




The Relationship Between Adult Attachment and Suicidal Thoughts and Behaviors: A Systematic Review

Tiago C. Zortea, Cindy M. Gray & Rory C. O'Connor


To cite this article: Tiago C. Zortea, Cindy M. Gray & Rory C. O'Connor (2019): The Relationship Between Adult Attachment and Suicidal Thoughts and Behaviors: A Systematic Review, Archives of Suicide Research

To link to this article: <https://doi.org/10.1080/13811118.2019.1661893>

 View supplementary material 

 Published online: 23 Sep 2019.

 Submit your article to this journal 

 View related articles 

 View Crossmark data 



The Relationship Between Adult Attachment and Suicidal Thoughts and Behaviors: A Systematic Review

Tiago C. Zortea , Cindy M. Gray , and Rory C. O'Connor 

We aimed to (1) determine the extent of the relationship between attachment and suicidal thoughts and behaviors (STB), (2) investigate whether any gender differences exist, and (3) conduct a theoretical synthesis of the included studies. A systematic search of the databases Web of Science; EBSCO Host Medical and Psychology-related resources, which includes CINAHL, Health Source: Nursing/Academic Edition, Psychology and Behavioral Sciences Collection, and PsycINFO; and EMBASE was conducted, with 52 studies meeting the inclusion criteria. While secure attachment acts as protective factor, insecure attachment orientations appear to be vulnerability factors compromising an individual's capacity to cope with relationship issues, increasing suicide risk. There is insufficient evidence about gender differences. The theoretical synthesis suggests that attachment, interacting with other psychological traits in response to negative life events, acts as a vulnerability or protective factor for STB. Implications for treatment are also discussed.

Keywords attachment, attachment theory, suicide, systematic review, suicidal ideation, relationships

INTRODUCTION

Globally, suicide is one of the three leading causes of death among people aged 15–44 years, and the second leading cause of death among young people aged 15–29 years (World Health Organization, 2014). There have been many advances in understanding which factors are associated with people's vulnerability to suicidal thoughts and behaviors (Hawton,

Saunders, & O'Connor, 2012; O'Connor & Nock, 2014; Turecki & Brent, 2016), including those related to family experiences (Brent et al., 1993; Cero & Sifers, 2013; Fergusson & Lynskey, 1995). *Suicidal thoughts and behaviors* (STB) is an umbrella term to describe suicide-related thoughts or ideation (including casual, transient, passive, active, and persistent thinking) and suicide-related behaviors—acts with some degree of suicidal intent,

including suicide attempts with and without injuries (Silverman, Berman, Sanddal, O'Carroll, & Joiner, 2007).

STB affects people of all ages, ethnicities, sexual orientation, and genders, although the overwhelming majority of suicide deaths are by men (World Health Organization, 2014). Although most studies on STB focus on the role of risk factors (Franklin et al., 2017), another approach to improving knowledge and prevention of suicide is by understanding better the motivations for suicide attempts (Klonsky, May, & Saffer, 2016), as well as how psychological and contextual variables interact to develop vulnerability for STB. Recent research has suggested that key elements associated with developmental vulnerability for STB are having been exposed to adverse childhood experiences (Cleare et al., 2018), harmful parenting (Cero & Sifers, 2013), and early disrupted attachment relationships (Fergusson, Woodward, & Horwood, 2000). Although several researchers have proposed a range of theoretical frameworks to understand the etiology of STB (Barzilay & Apter, 2014), Adam (1994) was the first author to suggest a developmental model where the attachment system plays a central role in the emergence of suicidal behavior. According to this model (Adam, 1994), early adverse parenting experiences and other childhood negative experiences could lead to the development of insecure attachments which could act as a psychological vulnerability factor for suicide risk later in life. Adam's model was based on Bowlby and Ainsworth's work which suggests a permanent effect of early attachment experiences (Ainsworth & Bowlby, 1991).

Through observational and experimental studies, Mary Ainsworth et al. (Ainsworth, Blehar, Waters, & Wall, 1978; Bretherton, 1992) proposed that the

attachment system functioned through different orientations (i.e., attachment styles or attachment patterns), which reflects the way internal working models of the self and others work. A secure attachment orientation is characterized by a self-perception of worthiness (lovability) connected with an expectation that other people are generally accepting and responsive. Insecure attachment may develop different orientations, anxious or avoidant, characterizing specific ways of emotional bonding with attachment figures (mothers or the closest carers in infancy, or partner and intimate relationships in adulthood). In general, anxious attachment (also known as preoccupied or ambivalent attachment) is characterized by an amalgamation of a sense of unworthiness (unlovability) and a positive appraisal of others, which leads the person to struggle for self-acceptance by gaining the acceptance of valued others. Finally, an avoidant attachment orientation includes behaviors that aim to produce and keep autonomy, control, and independence in their relationships, using "distancing" and "detaching" coping strategies, avoiding intimacy and emotional involvement. This is a result of having a negative view of their attachment figures and a negative view of the self (for those who present higher levels of attachment anxiety and attachment avoidance, "fearful-avoidant" or "disorganized" attachment patterns) or a negative view of their attachment figures and combined with positive view of the self (also known as "dismissive-avoidant" patterns) (Mikulincer & Shaver, 2008, 2016a; Simpson & Rholes, 2012).

In trying to understand the role of attachment in STB, it is important to consider how attachment is assessed. Due to the conceptual diversity within attachment theory, several authors have developed

different methods to measure adult attachment, including interviews and self-report measures (Ravitz, Maunder, Hunter, Sthankiya, & Lancee, 2010). These instruments are either based on typological approaches of attachment, which conceptualize secure, anxious, and avoidant relationships as categories (patterns or styles), or on dimensional approaches, which operationalize attachment across dimensions (levels) of anxiety and avoidance attachment. Although a typological approach seems to be useful in clinical settings (Ravitz et al., 2010), it has been criticized for losing individual variability, having poor reliability and validity properties, and lacking statistical power and precision. These measurement issues have been argued to be solved through dimensional approaches of assessing attachment orientations (Fralei, Heffernan, Vicary, & Brumbaugh, 2011; Fralei, Waller, & Brennan, 2000; Sibley, Fischer, & Liu, 2005).

Although Adam's (1994) theoretical framework suggests the existence of a relationship between attachment and STB, the nature of this relationship is still unclear, as is whether insecure attachment orientations (anxiety and avoidance) are specifically more associated with suicidal thoughts and/or suicidal attempts. Similarly, Adam's model does not propose distinct psychological and contextual factors related to attachment that could differentiate those who experience suicidal thoughts from people who attempt suicide.

Recent reviews have endeavored to summarize the available evidence between attachment and STB (Miniati, Callari, & Pini, 2017), and attachment and self-injurious behaviors (Wrath & Adams, 2018). Although the key message of the review by Miniati et al. (2017) is that insecure attachment style is associated with increased suicide risk, their review was not

systematically conducted. Wrath and Adams's (2018) review was systematic but its inclusion criteria were stricter, yielding only 17 studies in the final synthesis. For example, the latter review did not include studies assessing suicidal ideation. The present review not only extends the aims of the aforementioned reviews, but it also addresses some of methodological limitations of these reviews focusing on studies of STB (excluding non-suicidal self-injury and non-suicidal self-harm). In summary, the aims of this systematic review are to (1) investigate the extent of the relationship between attachment orientations and STB, including exploring potential mediators and moderators of the attachment and STB relationship, (2) investigate whether the relationship between attachment and STB is similar for men and women, and (3) provide a theoretical synthesis of the results presented by the included studies.

METHOD

Information Sources

A literature search was conducted using the following psychological and medical databases: Web of Science (all years, 1900–2017); EBSCO Host Medical and Psychology-related resources (all years, 1934–2017, which includes CINAHL, Health Source: Nursing/Academic Edition, Psychology and Behavioral Sciences Collection, and PsycINFO); and EMBASE (all years: 1946–2017).

Search (Full Electronic Strategy)

The search was conducted on March 6, 2017, with no date restriction. Keyword searches using the terms Attachment AND Suicid* were employed, which generated




FIGURE 1. Flow chart of the selection process.

2,389 records (1,003 after duplicates were eliminated). The search was refined by document types (articles) and languages (English; Figure 1).

Eligibility Criteria

Given that there were only 1,003 records, we took the decision to read all abstracts to determine whether they met the following inclusion criteria:

1. The study should be an empirical article.
2. The study should report a relationship between attachment and suicidal ideation or behavior.
3. The study should assess general attachment or attachment to parents.
4. The study should measure adults' or adolescents' perceptions of attachment or attachment patterns.

5. The study should be written in English.
6. The study should not be a single case study.
7. The study should not be a qualitative study.

Data Extraction and Quality Assessment

A data-extraction sheet was completed for each article with relevant information for this systematic review (e.g., article identification, methodological aspects, main results, and authors' interpretation of their data). After data extraction, all selected studies were assessed for methodological quality using a 9-item index based on O'Connor, Ferguson, Green, O'Carroll and O'Connor's (2016) quality assessment tool. Total scores were computed ranging from 0–13, meaning that a lower score is indicative of a higher probability of methodological bias. Classifications of quality were set as follows: 0–2 = *very low quality*, 3–4 = *low quality*, 5–7 = *reasonable/medium quality*, 8–10 = *good quality*, and 11–13 = *excellent/very good quality*. The detailed quality-assessment tool can be found in the [Supplementary material](#) (Table A). The checklist of items to include when reporting a systematic review or meta-analysis (Moher et al., 2009) was completed and can be found in the [Supplementary material](#) available (Table B).

Data Analysis and Synthesis

A narrative synthesis was conducted. The studies were grouped according to study design. An appraisal of authors' interpretations of their data was also conducted to identify psychological themes across the studies and compare these with predominant models of suicidal behavior. These models are the interpersonal theory

of suicide (IPT; Joiner, 2005), the integrated motivational-volitional model of suicidal behavior (IMV; O'Connor, 2011; O'Connor & Kirtley, 2018), the three-step theory (3ST, Klonsky & May, 2015), and the developmental model of suicidal behavior and attachment (Adam, 1994).

The first three contemporary models of suicidal behavior, known as ideation-to-action frameworks, propose that the factors associated with suicidal ideation (SI) are distinct from those associated with suicide attempts (SA). The IPT suggests that the combination of perceived burdensomeness and thwarted belongingness leads to suicidal thinking but the presence of acquired capability (e.g., fearlessness of death) is the key factor that distinguishes SI from SA. On the other hand, the IMV model suggests that defeat and entrapment are the main psychological factors that lead to SI, in which entrapment mediates the defeat-SI relationship. A range of volitional moderators, however, are hypothesized to differentiate between SI and SA. Finally, the 3ST suggests that SI arises from the combination of pain and hopelessness, and the SI-SA transition is facilitated by dispositional, acquired, and practical contributors to the capability for SA. Although none of the three new theories includes attachment explicitly, Adam's model (1994) suggests that attachment may act as a vulnerability or background variable acting as a general contributor to the development of STB.

RESULTS

Descriptive Overview of the Selected Studies, Quality Assessment, and Sample Characteristics

Of the 1,003 records identified, 52 were included in the systematic review (see [Figure 1](#)): 35 of the studies were

cross-sectional, 12 were prospective, three were case-control, one was a randomized controlled trial, and one used mixed methods (quantitative and qualitative analyses). Twenty-six (50%) were conducted in the United States; six in Canada; four in the United Kingdom; three in Portugal; two each in France, New Zealand, Turkey, and Italy; and one each in Israel, Iran, South Africa, South Korea, and Finland.

Although study quality varied, most studies were classified as reasonable/medium. The majority of the studies employed a cross-sectional design; therefore, it is not possible to comment about causality. The assessment of attachment and STB across the studies was heterogeneous, as the instruments were quite diverse (e.g., validated and non-validated self-report scales, interviews, single item/question, hospital admissions/records; references of the measures employed by the studies cited in Tables 1–3 can be found in the Supplemental material). Although most studies ($n = 43$) employed validated measures of adult attachment, only eight out of the 52 studies assessed the construct via clinical/research interviews. For the assessment of STB, 33 studies used self-administered measures and 19 employed interviews. Furthermore, 19 studies assessed STB through a non-validated scale or a single item/question; 10 used registers from hospital admission for suicide attempt or items from a validated diagnostic rating scale; and 23 assessed STB via clinical interview or a full validated scale.

There were 46,476 participants in the 52 studies (21,777 males and 24,689 females). The mean age was 25.1 years old ($SD = 11.8$) and 74% of participants ($n = 34,412$) were from the United States. Of the participants, 4,752 reported suicidal ideation or a suicide attempt and

25,893 did not report any STB. Fourteen studies did not provide details of the number of suicidal and non-suicidal participants. Two-thirds of participants (66.5%) were recruited from the general population, 16.1% were university students, 7.3% were clinical patients (inpatients, outpatients, and people looking for treatment or admitted to the hospital), 4.3% were a mix of clinical patients and participants from the general population, and 5.7% were participants from specific populations (those who are homeless, war veterans, or immigrants' children). In 24 studies, those in the suicidal group were clinically diagnosed with a mental health condition (nonpsychotic) ($n = 2,016$), whereas in two studies members of the suicidal group did not have any psychiatric diagnosis ($n = 406$). Twenty-six studies did not state whether members of the suicidal group had been diagnosed with a mental health condition ($n = 2,330$).

Associations Between Attachment and STB

The associations between attachment and suicidal ideation and behaviors are described in detail according to attachment patterns: secure attachment, avoidant attachment; anxious attachment; and other classifications of attachment.

Secure attachment. Thirty-three studies investigated the association between secure attachment and STB, and the vast majority of these associations were negative (i.e., higher levels of secure attachment were associated with lower levels of suicidal thoughts or suicide attempts). The following subsections show how secure attachment is related to suicidal ideation, suicide attempt, and STB.

Suicidal ideation. Twenty studies reported statistically significant associations

TABLE 1. Cross-Sectional Studies on Attachment and Suicidal Ideation/Behaviors.

Study no.	Population			Constructs and measures		Quality assessment
	Study/country	Sample size/source	Gender/age	Suicidal ideation/behaviors	Attachment	
1	Baiocco, Ioverno, Lonigro, Baumgartner, and Laghi, 2015. [Italy and Spain]	1,832, volunteers from the general population. 1,125 from Italy and 707 from Spain.	716 male, 1,116 female Mean age 25.66 (SD = 6.2)	<i>Construct:</i> SI. <i>Measure:</i> Single item from the Beck Depression Inventory (BDI-II; Beck, Steer, & Brown, 1996).	<i>Construct:</i> Attachment to peers. <i>Measure:</i> Short form of the Peer Subscale of the IPPA (Armsden & Greenberg, 1987).	2
2	Maimon & Kahl, 2008. [USA]	6,369, data from the National Longitudinal Study of Adolescent Health.	3,057 male, 3,312 female. Mean age 16.3 (SD = 1.62)	<i>Construct:</i> SA. <i>Measure:</i> Single question on having attempted suicide in the past 12 months.	<i>Construct:</i> AP (secure). <i>Measure:</i> Two non-validated questions on attachment to parents.	2
3	Nagra, et al., 2016 [UK]	323, people with current or past experience of self-harming.	38 male, 285 female. Mean age 22.86 (SD = 7.62).	<i>Construct:</i> S, SA. <i>Measure:</i> Suicidal Behaviors Questionnaire-Revised (SBQ-R; Osman et al., 2001).	<i>Construct:</i> AS. <i>Measure:</i> RQ (Bartholomew and Horowitz, 1991).	5
4	Palitsky et al., 2013. [USA]	5,692, data from the National Comorbidity Survey collected in 48 states of the U.S.	2,382 male, 3,310 female. Mean age 45 years old.	<i>Construct:</i> SI, SA. <i>Measure:</i> Self-reports of lifetime suicidal ideation and suicide attempts.	<i>Construct:</i> AS. <i>Measure:</i> Attachment self-report (Hazan & Shaver, 1987).	3
5	Peter et al., 2008. [Canada]	1,032, data from the National Longitudinal Survey of Children and Youth.	485 male, 547 female. Mean age 13.6 years old (SD = 1.1).	<i>Construct:</i> SI. <i>Measure:</i> Single question on having experienced suicidal ideation in the past 12 months.	<i>Construct:</i> AP (parental positive stimuli, parental negative stimuli, and parental involvement). <i>Measure:</i> 20-item non-validated scale.	2

(Continued)

TABLE 1. (Continued).

Study no.	Study/country	Population		Constructs and measures		Main results (effect sizes included when reported in the studies)	Quality assessment
		Sample size/source	Gender/age	Suicidal ideation/behaviors	Attachment		
<i>College and University Students</i>							
6	Buelow et al., 2000. [USA]	163, undergraduate students.	45 male, 109 female. Mean age 26 (SD = 7.6).	<i>Construct:</i> Suicidal thoughts and behaviors. <i>Measure:</i> The Suicide Behavior Questionnaire (Linehan & Nielson, 1981)	<i>Construct:</i> AS and object relations. <i>Measure:</i> The Attachment and Object Relations Inventory (Buelow, McClain, & McIntosh, 1996).	Secure attachment was an additional significant negative predictor of suicidal thoughts and behaviors ($F [2, 158] = 30.88$).	2
7	De Jong, 1992. [USA]	126, undergraduate students.	51 male, 75 female. Mean age 18.5.	<i>Construct:</i> S. (history of serious suicidal ideation or attempt). <i>Measure:</i> Non-validated self-report questionnaire.	<i>Construct:</i> APP. <i>Measure:</i> IPPA (Armsden & Greenberg, 1987).	Participants with history of SI and SA reported significantly lower levels of secure attachment to parents (mother: $d = 0.3714$; father: $d = 0.4295$) than control group.	2
8	Falgares et al., 2017. [Italy]	340, high-school students	92 male, 248 female. Mean age 16.5 (SD = 1.52)	<i>Construct:</i> S (lifetime suicidal ideation and attempts). <i>Measure:</i> Suicidal Behaviors Questionnaire-Revised (Osman et al., 2001).	<i>Construct:</i> AS. <i>Measure:</i> Attachment Style Questionnaire (Feeney, Noller, & Hanrahan, 1994)	Anxious attachment predicts suicidality mediated by self-criticism ($\beta = 0.15$) and dependency ($\beta = -0.05$). Avoidant attachment predicts suicidality mediated by self-criticism ($\beta = 0.13$).	6
9	Goncalves et al., 2016. [Portugal]	1,074, undergraduate students.	379 male, 695 female. Mean age 23.9 (SD = 6.10)	<i>Construct:</i> SI. <i>Measure:</i> Suicidal Ideation Questionnaire (Ferreira & Castela, 1999).	<i>Construct:</i> Attachment linkage prototypes. <i>Measure:</i> Adult Linkage Scale (Canavarro, Dias, & Lima, 2006)	Lower levels of comfort with proximity ($\beta = -0.10$) and trust in others ($\beta = -0.14$), and higher linkage anxiety ($\beta = -0.31$) were associated with higher SI risk.	3
10	Heydari et al., 2015. [Iran]	336, university students.	152 male, 184 female. Mean age 21.9 (SD = 2.38).	<i>Construct:</i> S (questions on suicidal ideation). <i>Measure:</i> 5-item Scale of Suicidality (Heydari et al., 2015).	<i>Construct:</i> APP. <i>Measure:</i> Attachment Scale (Ozbay & Ozcan, 2006) modified version for this study.	Suicidality correlated negatively significant with secure attachment (mother: $r = -0.33$; father: $r = -0.31$; Peer: $r = -0.35$). Secure maternal ($\beta = -0.14$) and peer attachment ($\beta = -0.16$) predicted suicidality via self-control and anomie.	0

11	Nunes & Mora, 2016. [Portugal]	604, high school, colleges, and university students.	274 male, 330 female. Mean age 15.99 (<i>SD</i> = 0.97)	<i>Construct:</i> SI. <i>Measure:</i> The Suicidal Ideation Questionnaire (SIQ; Reynolds, 1991)	<i>Construct:</i> AP. <i>Measure:</i> The Father/Mother Attachment Questionnaire (FMAQ; Matos & Costa, 2001).	Young people who indicate good quality of emotional bond to their parents reveal 19 times fewer SI ($\beta = 0.82$). Secure attachment to parents mediates the relationship between authoritative (father: $\beta = -0.19$; mother: $\beta =$ -0.17), authoritarian (father: $\beta = 0.05$; mother: $\beta = 0.06$) and permissive (father: $\beta = 0.03$; mother: $\beta = 0.04$) parenting styles, and SI.	4
12	Park et al., 2011. [South Korea]	249, university students.	66 male, 182 female. Mean age 21.8 (<i>SD</i> = 1.9).	<i>Construct:</i> suicide tendency. <i>Measure:</i> The Multi-Attitude Suicide Tendency (Orbach et al., 1991).	<i>Construct:</i> AAAS. <i>Measure:</i> ECRS (Brennan et al., 1998).	Levels of anxious attachment were positively correlated with levels of suicide proneness ($r = 0.15$).	2
13	Rodgers et al., 2011. [France]	615, high school students.	382 male, 233 female. Mean age 16.8 (<i>SD</i> = 1.3).	<i>Construct:</i> SI. <i>Measure:</i> Suicidal Behaviors Questionnaire-Revised (Osman et al., 2001)	<i>Construct:</i> AP (secure). <i>Measure:</i> IPPA (Armsden & Greenberg, 1987).	Attachment to parents was negatively correlated with SI (boys: $r = -0.27$; girls: $r = -0.23$). Attachment to parents predicted (negatively) SI mediated by defensive psychopathology and depression SI (boys $\chi^2 = 24.66$; girls: $\chi^2 = 10.99$, <i>ns</i>).	4
14	Strang & Orlofsky, 1990. [USA]	191, college students from two urban public universities.	92 male, 99 female. Mean age 19.5.	<i>Construct:</i> SI. <i>Measure:</i> Scale for Suicide Ideators (Schotte & Clum, 1982).	<i>Construct:</i> AP (secure) <i>Measure:</i> IPPA (Armsden & Greenberg, 1987).	The low SI group had less secure attachment than the non-SI group ($d =$ -0.53), and moderate to severe SI group had less secure attachment than non-SI ($d = -1.09$) and low SI groups ($d = -0.61$).	4
15	Zeyrek et al., 2009. [Turkey]	180, university students.	70 male, 110 female. Mean age 20.3 (<i>SD</i> = 1.6).	<i>Construct:</i> SI. <i>Measure:</i> The Suicide Probability Scale (Cull & Gill, 1982).	<i>Construct:</i> AS. <i>Measure:</i> RQ (Bartholomew & Horowitz, 1991; Scharf & Bartholomew, 1998).	SI was significantly correlated with secure (r $= -0.39$), preoccupied ($r = 0.29$), and fearful attachment ($r = 0.16$). Only for women, secure ($\beta = -0.18$), dismissing ($\beta = 0.19$), and preoccupied ($\beta = 0.18$) attachment patterns predicted variations in SI.	4

(Continued)

TABLE 1. (Continued).

General population		Population		Constructs and measures		Main results		Quality assessment
Study no.	Study/country	Sample size/source	Gender/age	Suicidal ideation/behaviors	Attachment	(effect sizes included when reported in the studies)		
<i>Patients</i>								
16	Aaltonen et al., 2016. [Finland]	287, patients from psychiatric units.	78 male, 209 female. Mean age 39.9 (SD = 13.0)	<i>Construct:</i> SI, SA. <i>Measure:</i> Questions adapted from the National Comorbidity Survey (NCS) (Kessler, Borges, & Walters, 1999).	<i>Construct:</i> AS. <i>Measure:</i> ECRS (Fraleigh et al., 2000).	Patients with history of repeated SA reported higher levels of anxious attachment ($d = 0.558$).	7	
17	Adam et al., 1996. [Canada]	133, patients (in- and out-) from psychiatric treatment centers.	74 male, 59 female. Mean age 15.3 (SD = 1.47).	<i>Construct:</i> S (defined as history of serious suicidal ideation or suicide attempt). <i>Measure:</i> Adam's (1973) Suicidal Ideation and Behavior protocol.	<i>Construct:</i> AS. <i>Measure:</i> AAI (George, Kaplan, & Main, 1985).	For the case group of adolescent patients with a history of STB, the predominant attachment pattern was unresolved-disorganized with preoccupied (41% of male patients and 30% of female patients in the case group). Women with disorganized attachment pattern and five or more other risk factors were 37 times more likely to belong to the suicidal/self-harm group (adjusting for age: AOR = 37.37).	5	
18	Bedi et al., 2014. [Canada]	167, treatment-seeking women who suffered from childhood abuse.	All female. Mean age 39.95 (SD = 11.11)	<i>Construct:</i> SH, SA. <i>Measure:</i> Structured interview adapted from the Major Depressive Episode section of the Dissociative Disorders Interview Schedule (Ross et al., 1989).	<i>Construct:</i> AS. <i>Measure:</i> Adult Attachment Projective System (George, West, & Pettem, 1997).	Low levels of secure attachment ($\eta^2 p^2 = 0.065$), high levels of fearful ($\eta^2 p^2 = 0.062$), and dismissive ($\eta^2 p^2 = 0.035$) attachment patterns are related to suicidal behavior in the presence of drug misusing.	4	
19	D'Orzio et al., 2015. [USA]	146, treatment-seeking African American women who attempted suicide in the prior year.	All female. Mean age 35.6 (SD = 11.43).	<i>Construct:</i> SA. <i>Measure:</i> Single self-report question on suicide attempt in the past 12 months.	<i>Construct:</i> AS. <i>Measure:</i> RSQ (Griffin & Bartholomev, 1994).	Attachment to mother and to peers ($\beta = -0.38$) independently predicted variations on SI. Attachment to mother and peers no longer accounted for significant variance in SI after adjusting for depressive symptoms.	5	

21	Gormley & McNeil, 2010. [USA]	109, short-term psychiatric inpatients from a university hospital.	65 male, 44 female. Mean age 40.7 (SD = 13.6). <i>Measure:</i> two questions on history of suicide attempt and non-suicidal self-injury. <i>Measure:</i> two questions on history of suicide attempt and non-suicidal self-injury since age 12.	<i>Construct:</i> self-directed aggression (history of suicide attempt and non-suicidal self-injury). <i>Measure:</i> RQ (Bartholomew & Horowitz, 1991).	Higher levels of attachment anxiety were significantly associated with history of SA (OR = 1.09).	5
22	Ilardi & Kaslow, 2009. [USA]	51, treatment-seeking African-American women reporting intimate partner violence and an SA.	All female. Mean age 34.5 (SD = 8.78). <i>Construct:</i> SA. <i>Measure:</i> Single self-report question on suicide attempt in the past 12 months.	<i>Construct:</i> AS. <i>Measure:</i> RSQ (Griffin & Bartholomew, 1994).	Participants were most likely to describe themselves as having a dismissive attachment pattern, followed by fearful and preoccupied attachment patterns.	3
23	Lessard & Moretti, 1998. [Canada]	116, clinical patients from a government-funded, provincial mental health facility.	73 male, 43 female. Mean age 13.4 (SD = 1.4). <i>Construct:</i> SI: (a) severity of suicidal wish and (b) lethality of method. <i>Measures:</i> (a) Youth Self Report of the Ontario Child Health Study Scales (Offord et al., 1987), and the BDI (Beck et al., 1961); and (b) Diagnostic Interview of Childhood and Adolescence-Revised (Reich, Shayka, & Tablisson, 1991).	<i>Construct:</i> AS. <i>Measure:</i> Semi-structured psychological intake interviews based on Bartholomew's (1990) attachment framework.	Youth with fearful attachment were 6.5 times more likely to endorse SI ($r = 0.42$). Elevated lethality of imagined method was correlated with higher ratings on the preoccupied pattern ($r = 0.59$).	5
24	Levi-Bez et al., 2013. [Israel]	102, consecutive patients after medically serious and non-serious suicide attempt at a university hospital.	49 male, 53 female. Mean age 37.3 (SD = 15.3). <i>Construct:</i> SI and degree of lethality of the SA. <i>Measures:</i> SIS (Beck, Schuyler, & Herman, 1974); The Lethality Rating Scale (Beck, Beck, & Kovacs, 1975).	<i>Construct:</i> AAAS. <i>Measure:</i> ECRS (Brennan et al., 1998).	Medically serious suicide-attempt patients showed higher levels of anxious attachment than non-serious suicide attempt patients ($d = 0.45$). Self-disclosure and loneliness mediated the paths between avoidant attachment and lethality of SA; loneliness mediated the paths between anxious and lethality of SA.	6
25	Lizardi et al., 2011. [USA]	524, depressed patients presenting for participation in research at a university hospital.	211 male, 313 female. Mean age 37 (SD = 13.3). <i>Construct:</i> SA. <i>Measure:</i> CUSHE (Oquendo, Halberstam, & Mann, 2003).	<i>Construct:</i> AS. <i>Measure:</i> AAS (Simpson, 1990).	The SA group showed higher levels of anxious attachment than the non-SA group ($d = 0.29$). Anxious attachment was associated with the likelihood of having made an SA (OR = 1.33).	7

(Continued)

TABLE 1. (Continued).

General population		Population			Constructs and measures		Main results (effect sizes included when reported in the studies)		Quality assessment
Study no.	Study/country	Sample size/source	Gender/age	Suicidal ideation/behaviors	Attachment				
26	Sheftall et al., 2013. [USA]	236, adolescents who have received inpatient psychiatric care.	124 male, 112 female. Mean age 14.48 years old.	Construct: SA. Measure: Suicidal Behavior Questionnaire (Addis & Linehan, 1989).	Construct: AP (secure). Measure: IPPA (Armsden & Greenberg, 1987).		Lower parental attachment levels for the SA group ($d=0.33$). Paternal attachment was the only significant predictor of SA status ($\beta = -0.014$).	6	
27	Sheftall et al., 2014. [USA]	80, presenting to behavioral health clinics or emergency department of a large children's hospital.	20 male, 60 female. Mean age 15.5 ($SD=1.35$).	Construct: SA. Measure: CUSHF (Maann et al., 1992).	Construct: AAAS. Measure: ECRS (Bremnan et al., 1998).		The SA group had higher levels of avoidant ($d=0.75$) and anxious attachment ($d=0.62$). Attachment avoidance was the only significant predictor of SA in the multivariable model ($OR = 2.05$).	7	
28	Stepp et al., 2008. [USA]	406, from psychiatric inpatient, outpatient, medical, and university settings.	136 male, 270 female, mean age 37.2 ($SD=10.5$).	Construct: Suicide-related behaviors. Measures: Part of standardized AXIS II diagnostic interviews for assessing one of the criteria for borderline personality disorder related to suicide and self-harm.	Construct: AAAS. Measure: A combination of factors from the ECRS (Fraleigh et al., 2000) and the Attachment Q-Sort (Kobak, 1989).		Anxious attachment increased the odds of belonging to the suicidal behavior group ($OR = 2.03$). Anxious attachment predicted suicidal behaviors mediated by interpersonal sensitivity ($\alpha = 2.05$), aggression ($\alpha = -2.29$), and lack of sociability ($\alpha = 2.04$). Avoidant attachment predicted suicidal behaviors mediated by interpersonal sensitivity ($\alpha = -1.99$).	9	
29	Venia et al., 2014. [USA]	114, adolescent inpatients admitted to an acute inpatient facility.	40 male, 74 female. Mean age 14.69 ($SD=1.47$).	Construct: SI. Measure: One item (number 9) from the BDI (Beck et al., 1996).	Construct: Attachment-related schemas Measure: Security Scale (Kerns et al., 2005).		Maternal attachment security is negatively correlated with suicide-related thoughts ($r = -0.351$). Secure attachment predicted variations in SI, mediated by thwarted belongingness ($\beta = -0.386$).	3	
30	West et al., 1999. [Canada]	187, adolescent inpatients admitted to a psychiatric treatment center.	102 male, 85 female. Mean age 14.9 ($SD = 1.6$).	Construct: S. Measure: The Suicidal Ideation and Behaviors Protocol (Adam, 1973).	Construct: Adolescent AS. Measure: The Adolescent Attachment Questionnaire (West et al., 1999).		Lower levels of secure ($\beta = 0.21$) and high levels of preoccupied attachment ($\beta = -1.64$) predict suicide risk.	6	

Specific Populations

31	Kidd & Shahar, 2008. [USA]	208, homeless who had no fixed address or were living in a shelter.	122 male, 86 female. Mean age 20.2 (SD = 2.39).	<i>Construct:</i> SI. <i>Measure:</i> Four items, Schedule for Affective Disorders and Schizophrenia for School-Age Children (Lewinsohn, Rohde, & Seeley, 1996).	<i>Construct:</i> AS. <i>Measure:</i> RQ (Bartholomew & Horowitz, 1991).	SI was positively correlated with fearful ($r = 0.23$) and preoccupied attachment ($r = 0.16$), and negatively with secured attachment ($r = -0.15$).	4
32	Nye et al., 2009. [USA]	48, treatment-seeking Vietnam combat veterans.	All male. Mean age 57.7 years old.	<i>Construct:</i> Current SI, lifetime SI. <i>Measure:</i> The Adam's Suicidal Ideation and Behaviors Protocol (Adam, 1973), & BSSI (Beck & Steer, 1993).	<i>Construct:</i> Attachment representation <i>Measure:</i> AAI (Main, Goldwyn, & Hesse, 2002).	The secure attached group demonstrated higher levels of SI ($d = 0.77$). The unresolved/disorganized group demonstrated lower levels of SI when controlling for PTSD severity ($d = 0.51$).	7
33	Rodell et al., 2003. [USA]	188, homeless veterans in a substance abuse program in a Veterans Affairs medical center.	178 male, 10 female. Mean age 44.5 (SD = 4.7).	<i>Construct:</i> SI. <i>Measure:</i> Hudson's (1990) Multi-Problem Screening Inventory.	<i>Construct:</i> levels of bonding (secure attachment) <i>Measure:</i> Hudson's (1990) Multi-Problem Screening Inventory.	Early positive attachment to caregivers (female caregiver: $\beta = -0.25$; male caregiver: $\beta = -0.27$) predicted a reduction of the probability of experiencing suicidal thoughts.	3
34	Van Leeuwen et al., 2010. [France]	292, high school immigrant adolescents.	170 male, 122 female. Mean age 17.2 (SD = 1.3).	<i>Construct:</i> SI. <i>Measure:</i> Suicidal Behaviors Questionnaire-Revised (Osman et al., 2001)	<i>Construct:</i> AP (secure). <i>Measure:</i> IPPA (Armsden & Greenberg, 1987).	Attachment to parents inversely predicted SI (boys: $\beta = -0.23$; girls: $\beta = -0.34$). The total score of attachment was negatively correlated with levels of SI ($r = -0.27$).	5

Patients and General Population

35	Pisetsky et al., 2015. [USA]	337, adult women from local eating disorder clinics and from the community	All female. Mean age 25.67 (SD = 8.85)	<i>Construct:</i> SA. <i>Measure:</i> Item on lifetime suicide attempt from the Impulsive Behavior Scale (Rosotto, Yager, & Rorty, 1994)	<i>Construct:</i> Insecure attachment <i>Measure:</i> Subscale of the Dimensional Assessment of Personality Pathology-Basic Questionnaire (DAPP-BQ; Livesley, Jang, & Vernon, 1998).	Insecure attachment predicted lifetime suicide attempt group membership (OR = 1.08). Insecure attachment predicted SA alongside identity problems, depression, and anxiety disorder ($\beta = 0.07$).	7
----	---------------------------------	--	---	---	---	---	---

Notes: SI: suicidal ideation; SA: suicide attempt; S: suicidality (authors' terminology for suicidal ideation and/or behaviors); SH: non-suicidal self-harm; BDI: Beck Depression Inventory; BHS: Beck Hopelessness Scale; AS: attachment style; AP: attachment to parents; APP: attachment to parents and peers; IPPA: Inventory of Parent and Peer Attachment; AAAS: avoidant and anxious attachment styles; ECRS: Experiences in Close Relationships Scale; RQ: The Relationship Questionnaire; AAI: Adult Attachment Interview; SIS: The Suicide Intent Scale; AAS: The Adult Attachment Scale; BSSI: Beck Scale for Suicide Ideation; RSQ: The Relationship Style Questionnaire; CUSHF: Columbia University Suicide History Form; PTSD: post-traumatic stress disorder.

TABLE 2. Prospective Studies on Attachment and Suicidal Ideation/Behaviors.

Study No.	Study/country	Population			Constructs, measures, and assessments/follow up		Quality assessment
		Sample size/source	Gender/age	Suicidal ideation/behaviors	Attachment	Main results (effect sizes included when reported in the studies)	
36	Fergusson et al., 2000. [New Zealand]	965, data were gathered over the course of a 21-year longitudinal Study of a birth cohort of 1265 children born in New Zealand.	479 male, 486 female.	Constructs: SI, SA. Measures: Questions on history of experiencing the constructs. Assessments: Conducted annually from ages 15–21.	Construct: AP (secure). Measure: IPPA (Armsden & Greenberg, 1987). Assessment: at age 15.	Attachment to parents predicted SI ($\beta = -0.04$) and SA ($\beta = -0.07$) along with family and personality variables. Participants reporting suicidal behavior were significantly more likely to come from families characterized by poor parent-child attachment and other factors.	8
37	Raudino et al., 2013 [New Zealand]	924, gathered as part of the Christchurch Health and Development Study from general population over 30 years.	449 male, 475 female.	Constructs: SI, SA. Measures: Questions on history of experiencing the constructs. Assessments: Conducted annually in two waves: from ages 21–25 and from 25–30.	Construct: AP (secure). Measure: IPPA (Armsden & Greenberg, 1987). Assessment: At age 15.	Increasing in parental attachment levels was correlated with declining rates of STB ($r = -0.11$). Alongside parental care and over-protection, low attachment levels predicted individuals' propensity to suicidal behaviors.	7
38	Riggs & Jacobvitz, 2002. [USA]	233, from 125 couples in the third trimester of a first-time pregnancy recruited through research ads.	115 male, 118 female. Mean age 30.5 years old.	Construct: SI. Measures: Self-report question on suicidal ideation. Assessments and follow up: Not informed.	Construct: AS. Measure: AAI (George et al., 1985). Assessments and follow up: Not informed.	Preoccupied adults more often reported SI than did secure or dismissing adults over time ($\chi^2[3] = 9.71$). Unresolved trauma (abuse) attachment patterns for women alone was related SI.	6
39	Turanovic & Pratt, 2015. [USA]	13,555, data from the National Longitudinal Study of Adolescent Health.	6,396 male, 7,162 female Mean age 22 years old.	Construct: S. Measures: Two dichotomous reports of suicidal thoughts and behaviors. Assessment: Conducted in the first second wave of interviews, participants aged 22.	Construct: Family attachment (secure). Measure: Six items on parents and family attachment. Assessment: Conducted in the first wave of interviews, participants aging 15.	Family attachment reduced the effects of STB on adult outcomes substantially for both males ($\beta = -0.09$) and females ($\beta = -0.10$).	9

Patients							
40	Chapman et al., 2009. [USA]	55, volunteers diagnosed with borderline personality disorder participating in an RCT at University of Washington.	All female, mean age 31.99 (SD=7.65).	<i>Constructs:</i> SI, SA, NSSI. <i>Measures:</i> The Suicide Attempt Self-Injury Interview (Linehan et al., 2006). <i>Assessments:</i> Conducted at pretreatment, 4-month intervals throughout the treatment, and 12-month follow up.	<i>Constructs:</i> Attachment (secure) <i>Measures:</i> Subscale of Reward Dependence-The Temperament and Character Inventory (Cloninger, 1992). <i>Assessments:</i> Conducted at pretreatment and post-treatment. <i>Constructs:</i> AS toward parents and peers.	Attachment was negatively correlated with SA ($r = -0.48$) and suicide intent ($r = -0.35$) over time. Lower levels of attachment predicted suicide intent at pre-Study time-point ($\beta = -0.32$). Lower levels of attachment predicted SA prospectively ($\beta = -0.30$).	7
41	Glazebrook et al., 2015. [UK]	52, adolescents referred to mental health services with recent history of self-harm.	3 male, 49 female. Mean age 15 years old.	<i>Construct:</i> Self-harm (SA + NSSI). <i>Measures:</i> The self-harm questionnaire (developed from the questions used in Hawton et al.'s (2012)). <i>Assessments:</i> 6-month follow up.	<i>Measures:</i> The Child Attachment Interview (Target, Fonagy, & Shmueli-Goetz, 2003); The Attachment Questionnaire for Children (Muris, Mayer, & Meesters, 2000). <i>Assessment:</i> 6-month follow up.	Participants who engaged in self-harm (including suicidal intent) at follow-up were classified as having insecure maternal attachment at baseline ($\chi^2[1] = 5.46$). Insecure maternal attachment (OR = 7.80) and poor peer attachment (OR = 8.01) independently predicted self-harm (including suicidal intent) at follow up.	5
42	Grunebaum et al., 2010. [USA]	135, patients who presented to the researcher's clinic institution for evaluation and treatment of major depressive episode.	55 male, 80 female. Mean age 39.2 (SD=12.3).	<i>Constructs:</i> SA, SI.	<i>Constructs:</i> AAAS.	Avoidant attachment independently predicted greater risk of SA during one-year follow-up of adults after presentation with a major depressive episode (HR = 1.16). Avoidant attachment predicted SI at three months (HR = 1.07).	10
43	Yaseen et al., 2014. [USA]	161, psychiatric patients hospitalized following suicidal ideation or suicide attempt.	76 male, 85 female. Mean age 37.76 (SD=13.0)	<i>Constructs:</i> SI, SA, and suicide triggers. <i>Measures:</i> Suicide Trigger Scale (Yaseen et al., 2014); Columbia Suicide-Severity Rating Scale (Posner et al., 2011); SIS (Beck, Steer, & Ranieri, 1988); <i>Assessments:</i> 6-month follow up.	<i>Construct:</i> AS. <i>Measures:</i> RSQ (Griffin & Bartholomew, 1994). <i>Assessments:</i> 6-month follow up.	Participants with ultra-high levels of suicide risk reported significantly higher scores of fearful attachment. Participants with ultra-low levels of suicide risk reported significantly higher scores of secure attachment pattern.	10

(Continued)

TABLE 2. (Continued).

Study No.	Study/country	Population			Constructs, measures, and assessments/follow up		Main results (effect sizes included when reported in the studies)	Quality assessment
		Sample size/source	Gender/age	Suicidal ideation/behaviors	Attachment			
<i>School and University Students</i>								
44	Gilreath et al., 2009. [South Africa]	657; data from South Africa epidemiology high school survey.	296 male, 361 female. Mean age 14 (SD = 1.1).	<i>Construct:</i> SI. <i>Measure:</i> Two self-report questions on suicide thoughts in the past 12 months. <i>Assessments:</i> 2-year follow up.	<i>Construct:</i> Maternal closeness (secure attachment to mother). <i>Measure:</i> Two self-report questions on participant-mother closeness. <i>Assessment:</i> 2-year follow up.	Among females, maternal closeness was a negatively significant predictor of SI ($\beta = -0.145$). SI was a significant mediator of the effect of maternal closeness on Lifetime Smoking ($\beta = 0.381$).	3	
45	Liu, 2005. [USA]	2,966; Data from the National Longitudinal Study of Adolescent Health with school based design.	1,426 male, 1,540 female. Mean age 15 years old.	<i>Construct:</i> SI. <i>Measure:</i> Single self-report questions on suicide thoughts in the past 12 months. <i>Assessments:</i> 1-year follow-up.	<i>Construct:</i> Closeness with parents (secure AP). <i>Measure:</i> 10-item self-report scale on participant-parents closeness. <i>Assessments:</i> 1-year follow-up.	For girls, attachment to fathers independently predicted reduction of SI over time ($\beta = -0.24$). For boys, attachment to mothers independently predicted reduction of SI over time ($\beta = -0.24$).	6	
<i>Patients and General Population</i>								
46	Salzinger et al., 2007. [USA]	153; patients (New York City Register for Child Maltreatment) and control from general population.	93 male, 60 female. Mean age 16.5 (SD = 0.53).	<i>Construct:</i> SI, SA. <i>Measure:</i> Youth Risk Behavior Survey (Garrison, McKeown, Valois, & Vincent, 1993). <i>Assessment:</i> Conducted at time 2 (at participants' 16.5 years of age).	<i>Construct:</i> AP (secure). <i>Measure:</i> IPPA (Armsden & Greenberg, 1987). <i>Assessment:</i> Conducted at participants' ages 10.5 (time 1) and 16.5 years old (time 2).	Attachment to parents when participants were 16.5 years old independently decreased risk of SI (OR = 0.43) and SA (OR = 0.39).	9	
<i>Specific Populations</i>								
47	Smith et al., 2016. [USA]	2107, U.S. military veterans.	1,919 male, 188 female. Mean age 60.3 (SD = 15.0)	<i>Construct:</i> SI. <i>Measure:</i> Modified item from the Patient Health Questionnaire-9 (Kroenke & Spitzer, 2002).	<i>Construct:</i> AS. <i>Assessment:</i> Question assessing feeling and attitudes in relationships (Hazan and Shaver, 1994).	Lower levels of secure attachment predicted SI over time (two years later) (RR = 4.37).	12	

Notes: SI: suicidal ideation; SA: suicide attempt; S: suicidality (authors' terminology for suicidal ideation and/or behaviors); NSSI: non-suicidal self-injury; AS: attachment style; AP: attachment to parents; IPPA: Inventory of Parent and Peer Attachment; AAAS: avoidant and anxious attachment styles; AAI: Adult Attachment Interview; SIS: The Suicide Intent Scale; BSSI: Beck Scale for Suicide Ideation; RSQ: The Relationship Style Questionnaire; CUSHF: Columbia University Suicide History Form.

TABLE 3. Other Study Designs on Attachment and Suicidal Ideation/behaviors

Study No.	Study/country	Population		Constructs and measures		Main results (effect sizes included when reported in the studies)	Quality assessment
		Sample size/source	Gender/age	Suicidal ideation/behaviors	Attachment		
48	Cruz et al., 2013. [Portugal]	1,308, adolescents from community (control = 1,266) and patients with suicidal behaviors (clinical sample = 42).	609 male, 699 female. Mean age = 15.87 (SD=2.11).	<i>Constructs:</i> History of self-destructive thoughts and behaviors. <i>Measure:</i> Questions on suicidality from The Youth Self-Report (Achenbach, 1991)	<i>Construct:</i> AP (Secure) <i>Measure:</i> The Father/Mother Attachment Questionnaire (Matos, Barbosa, & Costa, 2001)	Being female, perceiving poor attachment to parents, a low level of control and a high level of rejection from one's father (OR = 0.58), a high degree of control from one's mother (OR = 0.003), and a low level of family cohesion increase the probability of membership in the self-destructive thoughts and behaviors group.	6
49	Özer et al., 2015. [Turkey]	122, psychiatric patients and controls from the general population.	20 male, 109 female. Mean age = 31.48 (SD=9.71).	<i>Constructs:</i> SI, SA. <i>Measures:</i> Scale of Suicidal Ideation (Dilbaz, Holat, Bayram, & Tüzer, 1995); Suicidal Behavior Scale (Bayram, Dilbaz, Bitlis, Holat, & Tüzer, 1995)	<i>Construct:</i> AAAS. <i>Measure:</i> ECRS (Brennan et al., 1998).	SA and depressive groups reported higher levels of avoidant ($d = -0.07$) and anxious ($r = 0.40$) attachment. The rate of patients with SA was found to be higher in the group of patients with the fearful attachment style ($r = 0.51$).	5
50	Violo & Arato, 2004. [Canada]	52, suicidal clinical adolescents ($n = 17$) and community control group ($n = 35$).	24 male, 28 female. Mean age 14.7 (SD = 1.7)	<i>Construct:</i> SA. <i>Measure:</i> Clinical diagnostic based on the <i>Diagnostic Statistical Manual</i> (APA, 2000).	<i>Construct:</i> Levels of secure attachment. <i>Measure:</i> Adolescent Attachment Survey (Genuis, Violano, Oddone-Paolucci, Robitaille, & McBride, 2000).	The SA group had significantly higher levels of insecure parental attachment ($r = 0.52$).	5
<i>Randomized Controlled Trial</i>							
51	Rossouw & Fonagy, 2012. [UK]	80, adolescent psychiatric patients with self-harm (inclusive suicidality).	12 male, 68 female. Mean age 14.7.	<i>Construct:</i> Self-harm (suicidality inclusive) <i>Measure:</i> Risk-Taking and Self-Harm Inventory (Vrouva, Fonagy, Fearon, & Rousow, 2010).	<i>Construct:</i> AAAS. <i>Measure:</i> ECRS (Brennan et al. 1998).	Avoidant attachment style independently predicted variations of self-harm (including suicidal intent) at the end of the treatment ($\beta = -0.17$). Avoidant Attachment mediates the relationship between Mentalization-Based Treatment for Adolescents and variances in Self-harm/SA ($\beta = -5.95$).	9

(Continued)

TABLE 3. (Continued).

Study No.	Study/country	Population			Constructs and measures		Main results (effect sizes included when reported in the studies)	Quality assessment
		Sample size/source	Gender/age	Suicidal ideation/behaviors	Attachment			
<i>Mixed Methods Study</i>								
52	Wright et al., 2005, [UK]	35, clinical population undergoing assessment for weekly psychotherapy. Control group from general public.	12 male, 23 female. Mean age 17.	Construct: Suicide risk. Measure: Pfeiffer Child Suicide Scale (Pfeiffer, 1986).	Construct: Separation anxiety. Measure: The Adolescent Separation Anxiety Interview (Richard, Fonagy, Smith, Wright, & Binney, 1998).	The high suicide risk group included a majority of adolescents who gave preoccupied attachment narratives and no secure narratives. The low suicide risk group was more diversely spread across the three main insecure attachment patterns, and the control sample over-represented the secure category.	4	

Notes: SA: suicide attempt; AP: attachment to parents; AS: attachment style; AAAS: avoidant and anxious attachment styles; ECRS: Experiences in Close Relationships Scale.

between secure attachment and suicidal ideation (studies 1, 10, 11, 13–15, 20, 23, 29, 31–34, 36, 39, 40, and 44–47; main results are in Tables 1–3). With the exception of Study 32, all studies found secure attachment to be a protective factor against suicidal ideation. Study 32, which explored suicidal ideation among Vietnam combat veterans in treatment for PTSD, found the secure attachment group had higher levels of current suicidal ideation than the insecure attachment group: however, given the relatively small sample size ($N=48$) and specific population, these results should be interpreted cautiously. In nine studies, suicidal ideation was negatively correlated with secure attachment (studies 10, 11, 13, 15, 23, 29, 31, 34, and 40). Other studies conducted multivariate analyses controlling for a range of variables alongside secure attachment to explain the variance in suicidal ideation (10, 11, 13, 15, 20, 29, 33, 34, 36, 39, 40, and 44–47). Within these studies, secure attachment predicted a reduction in suicidal ideation even when confounders were controlled for.

Suicide attempt. All 10 studies found a negative relationship between secure attachment and suicide attempt (studies 2, 3, 4, 19, 26, 36, 39, 40, 41, and 46). In particular: (a) high levels of secure attachment were negatively correlated with frequency of suicide attempts (study 40); (b) suicide-attempt survivors reported significantly lower levels of secure attachment compared with those who have not attempted suicide (studies 3 and 26); (c) high levels of secure attachment predicted lower frequencies of suicide attempts (studies 40, 41, 2, 4, and 26); and (d) low levels of secure attachment predicted suicide attempt (studies 36, 39, and 46). The relationship appears robust as some studies (2, 4, 39, 40, 41, and 46) controlled for

confounders (e.g., sociodemographic factors, depression, self-esteem, religiosity), and only one (study 40) reported a non-significant association between secure attachment and current suicide attempt (after adjusting for previous suicide attempt). Study 19 compared a “drug misuse problem” group with a “no drug misuse problem” group, with all participants having attempted suicide in the previous year; the former group showed lower levels of secure attachment. Study 3 included individuals with current and past experiences of self-harm recruited from an online community, and study 26 included adolescents from an inpatient psychiatric unit without previous self-harm.

STB. Eight studies did not differentiate between suicidal ideation and suicide attempt, but referred to “suicidality” as a mixed concept (i.e., including STB). They all reported a negative relationship between secure attachment and STB (studies 3, 6, 7, 30, 37, 43, 48, and 50). In particular, (a) there was a negative correlation between secure attachment and STB (studies 3 and 37); (b) suicidal groups reported significantly lower levels of secure attachment (studies 7, 43, and 50); and (c) lower levels of secure attachment predicted STB alongside other psychosocial variables in multivariate analyses (studies 3, 6, 30, 37, and 48). In the three studies (studies 30, 37, and 43) that controlled for covariates, the association remained significant after statistical adjustment.

The vast majority of the studies that investigated the relationship between secure attachment and STB found that secure attachment was protective against STB. Even when considering the methodological heterogeneity across the studies, this relationship appears to be robust.

Anxious attachment. 18 studies assessed anxious attachment and how it is

associated with STB. This section includes both anxious and preoccupied attachment because they address the same psychological characteristics of attachment anxiety. It is important to note that attachment anxiety and anxious attachment are used synonymously across the studies.

Suicidal ideation. Seven studies investigated the relationship between anxious or preoccupied attachment and suicidal ideation (studies 12, 15, 23, 28, 31, 38, and 52). Higher levels of attachment anxiety were positively correlated with suicidal ideation (studies 12, 15, 23, and 31), and suicidal ideation groups reported higher levels of attachment anxiety when compared to controls (studies 23 and 52). One study (study 38) comparing levels of suicidal ideation across attachment pattern groups (secure, anxious and avoidant) reported that the anxious group had higher levels of suicide ideation than the other groups. Study 15 showed that preoccupied attachment (together with hopelessness and poor problem-solving skills) explained variance in suicidal ideation. Study 28 presented findings indicating that attachment anxiety predicted membership of the suicidal ideation group and this relationship was mediated by interpersonal problems. This was the only study to investigate the relationship between suicidal ideation and anxious attachment that controlled for age, ethnicity, and gender, and demonstrated significant associations after adjustment.

Suicide attempt. Nine studies investigated the relationship between anxious attachment and suicide attempts (studies 3, 16, 21, 22, 24, 25, 27, 28, and 49). One study (study 24) found a positive correlation between the lethality of suicide attempts and the level of anxious attachment (the higher levels of anxious attachment, the more lethal a suicide attempt

was). The remainder found that patients who had made a medically serious suicide attempt showed higher levels of anxious attachment than patients who were deemed non-medically serious (medically serious suicide attempts were defined as more than 24 hours of hospitalization in an intensive care unit or surgery under general anesthesia after a suicide attempt). Study 16 found that patients with two or more lifetime suicide attempt presented higher levels of anxious attachment when compared to other groups of patients (no suicidal ideation group, suicidal ideation with no attempts group, and one suicide attempt group) after controlling for traits of borderline personality disorder.

Four studies using multivariate analyses reported that attachment was a significant predictor of suicide attempt: (a) study 21 found that depressive symptoms mediated the relationship between attachment anxiety and suicide attempt; (b) study 24 showed that attachment anxiety, alongside depression, hopelessness, and negative life events, predicted high levels of lethality in suicide attempts and that avoidant attachment explained the variance in medical lethality of suicide attempt mediated by self-disclosure and loneliness; (c) study 25 found that age and anxious attachment predicted the likelihood of having made a suicide attempt. Finally, (d) study 28 conducted a mediation analysis and demonstrated that attachment anxiety predicted suicide attempt group membership, but this relationship was mediated by lack of sociability. Other studies (16, 21, 27, and 28) controlled for covariates (e.g., gender, age, ethnicity, borderline personality disorder traits, depressive symptoms), and all reported significant associations between anxious attachment and suicide attempt.

STB. Three studies (studies 8, 17, and 30) explored the association between

anxious attachment pattern and STB. Study 8 found that anxious attachment explained the variance in STB through the mediation of dependency and self-criticism, after adjusting for gender. Study 17, assessing teenagers in psychiatric treatment, showed that those who were female, older than 15 years, and had unresolved-disorganized attachment-related trauma and an anxious attachment pattern were significantly more likely demonstrate STB. Finally, study 30 found that being older and having an anxious attachment pattern increased the likelihood of belonging to the STB group, after controlling for age and gender. Having an anxious (or preoccupied) relationship orientation seems to be a vulnerability factor for both suicidal ideation and suicide attempt even when other psychological variables are controlled for.

Avoidant attachment. Seventeen studies looked at the relationship between avoidant attachment orientation and its association with suicidal thoughts and actions. This section includes studies reporting on avoidant attachment orientations in general, as well as the dismissive avoidant and fearful avoidant patterns. Attachment avoidance and avoidant attachment are used interchangeably across the studies.

Suicidal ideation. Suicidal ideation was associated with (a) avoidant attachment (studies 4 and 42), (b) fearful avoidant attachment (studies 15, 23, and 31), and (c) dismissive avoidant attachment (studies 15 and 23). Specifically, avoidant attachment predicted variance in suicidal ideation at two time points (baseline and 3-month follow up; study 42), and suicidal ideation group membership in study 4 (when adjusted for several sociodemographic and mental health variables). Levels of fearful avoidant attachment were

positively correlated with suicidal ideation (studies 15, 23, and 31), and participants within the suicidal ideation group reported higher levels of fearful avoidant attachment than controls (study 23). The picture in relation to dismissive avoidant attachment is less clear: Study 15 found a positive correlation between levels of dismissive avoidant attachment and suicidal ideation but only among female participants, and a negative correlation was found between dismissive avoidant attachment and suicidal ideation in study 23.

Suicide attempt. Most of the studies that investigated the relationship between avoidant attachment and STB found it to be more strongly associated with suicide attempts than suicidal thoughts (studies 3, 4, 16, 19, 22, 24, 27, 28, 42, 49, and 51). In particular, (a) levels of avoidant attachment were positively correlated with levels of medical lethality of suicide attempts (study 24), and (b) higher levels of attachment avoidance among those who attempted suicide in the past compared to those who had never engaged in suicidal behavior or had suicidal ideation (study 27). Six studies (4, 24, 27, 28, 42, and 51) investigated avoidant attachment as part of a multivariate model predicting suicide attempt. Study 28 found that interpersonal sensitivity mediated the relationship between avoidant attachment and the risk of suicide attempt. Three studies (3, 19, and 22) investigating the relationship between dismissive and fearful avoidant attachments and suicide attempt showed that: (a) people with a history of suicide attempts reported higher levels of dismissive and fearful avoidant attachment than those with no history of suicide attempts (study 3); (b) women with a history of suicide attempt and drug misuse problems had significantly higher dismissive and fearful attachment scores than those with

no history of drug misuse (study 19), and (c) participants with a history of suicide attempt were most likely to describe themselves as having a dismissive and avoidant attachment pattern in comparison with other attachment patterns (study 22). Study 49 found higher levels of fearful avoidant attachment in people with a history of suicide attempts compared with patients with major depression and health controls. Of the eleven studies that focused on the relationship between suicide attempt and avoidant attachment orientations, only six (4, 16, 27, 28, 42, and 51) controlled for confounders and found no post-adjustment changes in their results.

STB. Four studies found associations between avoidant attachment orientations and STB (3, 8, 17, and 43). Study 8 found that avoidant attachment predicted variation in STB, and this relationship was mediated by self-criticism. Two studies (3 and 17) found dismissive avoidant attachment and STB to be associated but in different ways: Study 3 revealed a positive correlation between levels of dismissive avoidant attachment and STB, even after controlling for other psychological variables, but study 17 found that dismissive avoidant attachment was associated with the absence of suicidal ideation or attempt. Authors of this latter study suggested that it is unlikely that suicidal feelings were simply denied in the dismissive avoidant group and, therefore, future longitudinal research was needed. Studies 3 and 43 found that (a) levels of fearful avoidant attachment were positively correlated with levels of STB (study 3), and (b) participants with ultra-high levels of suicide risk reported significantly higher scores of fearful avoidant attachment after controlling for confounders (study 43).

Consistent with the pattern of findings found in the studies reporting an anxious

attachment–STB relationship, the literature also suggests that an avoidant orientation of attachment acts as a vulnerability factor for STB in interaction with other psychological variables. In most of the studies, this finding also appears to hold regardless of how the type of avoidant attachment is conceptualized (i.e., dismissive or fearful).

Other Attachment Classifications.

The conceptualization of attachment patterns may vary across theoretical views on attachment in the literature. Although Ainsworth's original proposition established three basic attachment patterns (secure, anxious/ambivalent, or avoidant), attachment is used as a synonym for a diverse range of definitions across the studies, including emotional bonding (33), object relations (6), parental involvement (5), parental closeness (44), and separation anxiety (52). Therefore, some studies have assessed attachment differently from the more frequently used classifications (secure, anxious, and avoidant). In this section, we describe the results of those studies that adopt other conceptual attachment patterns and their associations with suicidal thoughts and actions.

Suicidal ideation. Two studies (32 and 38) assessed unresolved/disorganized attachment. In Study 38, authors define unresolved attachment as a state of mind, “not an enduring, organized attachment strategy but rather a brief collapse of an adult's mental organization during discussions of trauma” (p. 196). The authors found that unresolved adults (especially females) were significantly more likely than other attachment groups to report emotional distress and suicidal ideation. In study 32, those in the unresolved/disorganized group demonstrated lower levels of current suicidal ideation when controlling for PTSD severity than those in the secure

group. These results should be interpreted cautiously, though, given the relatively small sample size ($N=48$). Moreover, the findings may not be generalized beyond the existing sample of Vietnam combat veterans in treatment for PTSD.

In study 5, secure and insecure attachment were assessed as “parental positive stimuli” and “parental negative stimuli,” respectively, with parental negative stimuli being the only predictor of suicidal ideation group membership. Finally, in study 9, attachment patterns were assessed via “linkage prototypes” (defined as relationship proximity patterns). Using this operationalization of attachment, suicidal ideation was found to be associated with lower levels of “comfort with proximity” and “trust in others” and with higher levels of linkage anxiety.

Suicide attempt. Study 18 assessed attachment through a projective test comprising seven black-and-white images of people illustrated in several circumstances eliciting themes of separation, loneliness, and loss. The vast majority of the participants with a history of suicide attempt were classified as having unresolved attachment (73.7%, $n = 123$). Eleven were classified with preoccupied attachment pattern (6.6%), 29 with dismissing attachment pattern (17.4%), and three (1.7%) were classified as secure. In this study, participants with a disorganized attachment pattern and five or more risk factors were 37 times more likely to belong to the suicidal group than those with zero risk factors. Study 41 employed a longitudinal design to assess attachment to parents among adolescents who had engaged in suicide attempt. It found that insecure maternal attachment independently predicted suicide attempt at 6-month follow up. Finally, study 35 evaluated general insecure attachment among women diagnosed

with bulimia nervosa, with 27.5% of them reporting a suicide attempt history. These authors found that: (1) the lifetime suicide attempt group had significantly higher levels of insecure attachment than the no suicide attempt group; (2) insecure attachment predicted lifetime suicide attempt group membership (univariate logistical regression); and (3) insecure attachment alongside identity problems, depression, and anxiety disorder predicted the suicide attempt group membership (multiple logistic regression).

STB. Only study 17 assessed the relationship between different classifications for insecure attachment and STB. The study reported that adolescents who were female, older than 15, and had unresolved-disorganized attachment-related trauma and preoccupied attachment pattern had a significant increase in the probability of membership in the STB group. By unresolved-disorganized attachment-related trauma the authors meant a lack of resolution of traumatic events related to attachment such as the death of a parent or other close relationship figure or the lack of resolution of physical or sexual abuse. In this study, 77% of all participants classified as having unresolved-disorganized attachment-related trauma or preoccupied attachment pattern were in the suicidal group.

Moderating and Mediating Factors

Out of the 52 studies of the current review, only eight (8, 10, 13, 21, 24, 28, 29, and 45) investigated the effect of mediators or moderators of the attachment-STB relationship. Three studies (10, 24, 28, and 29) found significant associations for the mediating effect of interpersonal factors: self-disclosure and loneliness (24), interpersonal sensitivity, aggression, and

lack of sociability (28), anomie (broken social bonds, study 10), and thwarted belongingness (29). Studies 8 and 10 reported evidence for the mediating effects of intrapersonal characteristics: self-criticism (8) and self-control (10). Two studies (13, 21) tested the role of psychiatric symptoms as mediators between attachment and STB: both studies found significant associations for depressive symptoms, and study 13 also found evidence for the mediating effect of defensive psychopathology, substance use, borderline features, and dissociative symptoms. Only one study (study 45) conducted a moderator analysis and found moderating roles for demographic characteristics (age and ethnicity).

Gender Differences

Only seven out of the 52 studies within this systematic review (14, 17, 18, 38, 44, 45, and 48) reported on gender differences or effects specific to males or females. Four studies (19, 22, 35, and 40) recruited only female participants. The studies that reported gender differences, with the exception of study 45, showed that insecure attachment increased suicide risk if the participants were female.

Studies 14 and 44 reported that the association between secure attachment and suicidal ideation was significant only for female participants. Study 45 found that for girls, attachment to fathers independently predicted reduction of suicidal ideation over time and for boys, attachment to mothers independently predicted reduction of suicidal ideation over time. One study (study 48) reported a gender effect increasing suicide risk: Being female, perceiving poor attachment to parents, a high level of paternal rejection, a high degree of maternal control, and a low level of family

cohesion increase the probability of membership in the self-destructive thoughts and behaviors group.

Although there was some evidence of a potential gender effect for the attachment-STB relationship, the current evidence is inconclusive as there was no consistent pattern of findings across the studies and often the gender analyses appeared to be post hoc.

Theoretical Explanations of the Relationship between Attachment and STB

Although the studies yielded evidence of the existence of an association, an important question to be addressed is *why* attachment and STB are associated. Taking an overview of these studies, most authors argued that, in interaction with other personality factors, attachment moderates or mediates the relationship between current negative events and psychological states that, in turn, may lead to STB. More specifically, (1) current negative events interpreted as threatening contextual cues would (2) activate the attachment system which, through one's models of the self and of others, would interact with other personality traits to act as coping strategies to deal with those negative events and with the emotional reactions to those events. This interaction would (3) trigger certain psychological states if those strategies fail to work, which would subsequently lead to (4) STB when a resolution of those current negative events is not met.

An example could be: (1) in contexts of a high likelihood of being criticized by others, such as an experience of being bullied, (2) individuals with higher levels of anxious attachment generally develop a negative internal working model of the self and tend to engage in self-critical verbal

behavior and harsh self-evaluation (and other traits of neuroticism and self-criticism) aimed at reducing the likelihood of being criticized by others. If this strategy fails to work, (3) other psychological states such as depression, hopelessness, and isolation may ensue and lead to (4) STB if the experiences of being bullied remain.

DISCUSSION

This systematic review set out to examine the relationship between attachment and STB in adolescents and adults. A key finding was that secure attachment functions as a protective factor against STB, and insecure attachment (variations of avoidant and anxious attachment orientations) seems to increase risk for suicidal thinking and behavior. Although the literature clearly shows the existence of an association between attachment and STB, the precise nature of the relationship between insecure attachment orientations and STB remains unclear; in the vast majority of the studies, all forms of insecure attachment orientations were found to be associated with both suicidal ideation and suicide attempt. This suggests that it is not possible to determine whether a specific orientation or style of insecure attachment differentiates between suicidal ideation and suicide attempt groups. As a consequence, it is difficult to identify a single explanation for why anxious attachment, for instance, would be positively associated with suicidal ideation but not with suicide attempts. Wrath and Adams's (2018) review reported that attachment anxiety is a crucial factor for engaging in both forms of self-injurious behavior (non-suicidal self-harm and suicide attempt), but did not demonstrate whether specific forms of insecure attachment differentiated between

those who engaged in self-injurious behavior from those who attempted suicide. Despite including a wider range of studies, our review yielded similar findings in that we found it difficult to determine whether anxious and avoidant attachment orientations were differentially associated with histories of suicidal ideation versus suicide attempt.

A secondary aim was to investigate whether there are gender differences in the relationship between attachment and STB. Although gender differences regarding suicide are prominent and well documented in suicide research more generally, the same cannot be concluded for the attachment-STB literature. Few studies reported the effect of gender on the relationship between attachment and STB. Although six studies demonstrated that insecure attachment increased suicide risk in women, the generalizability of these findings should be interpreted with caution because the majority of the studies were cross-sectional and they represent only 21% of the total articles in this systematic review and 12.2% of the total number of participants. Moreover, it was often difficult to investigate the relationship in men because there were too few male participants in the majority of these studies. Thus, there is insufficient evidence to draw any conclusions on gender differences on the relationship between attachment and STB.

Regarding factors that may elucidate our understanding of the relationship between adult attachment and STB, only eight out of the 52 studies (15.3%) tested the role of interpersonal, intrapersonal, psychopathological, and demographic factors. Although the inclusion of such factors was theoretically justified by the studies' authors, only one study (29) tested a component of a contemporary theory of

suicidal behavior (IPT; thwarted belongingness; Joiner, 2005). The attachment-theory research field would benefit from investigating the extent to which psychological variables suggested by the ideation-to-action models of suicidal behavior help understand the attachment-STB relationship (Joiner, 2005; Klonsky & May, 2015; O'Connor, 2011; O'Connor & Kirtley, 2018).

Nonetheless, it is possible to note that the relationship between attachment and STB is consistent with contemporary frameworks of suicidal behavior. For example, Adam's (1994) model of suicidal behavior, focused on how the vulnerability for STB develops across the lifespan, posits that attachment insecurities in childhood may act as distal risk factors for STB later in life (e.g., studies 11, 36, and 37). Although the vast majority of the studies in this review are not longitudinal, their findings are consistent with Adam's hypothesis of attachment serving as an intermediate risk factor connected to other psychological states and personality traits preceding STB.

The theoretical synthesis also suggests that types of insecure attachment may act as a diathesis as described in the integrated motivational-volitional model of suicidal behavior (O'Connor, 2011). According to the latter, in conjunction with environmental influences, life events, and other personality traits (e.g., social perfectionism, loneliness), forms of insecure attachment may act as a psychological vulnerability for STB. Insecure attachment orientations may increase one's sensitivity to feeling defeated and trapped (psychological states), thereby elevating the likelihood that suicidal thoughts emerge. Likewise, insecure attachment may also facilitate the development of thwarted belongingness and burdensomeness proposed in the interpersonal

model of suicide (Joiner, 2005). In Study 29, for example, Venta et al. (2014) suggest that interpersonal risk factors for STB can potentially be targeted by addressing beliefs of thwarted belongingness, rather than treating attachment-related risk for STB as static. Moreover, activation of the attachment system could also increase emotional pain and weaken feelings of connectedness as outlined in the three-step theory (Klonsky & May, 2015), increasing the risk of developing STB. Although most authors mentioned the potential interaction of attachment with proximal psychological and interpersonal factors to increase suicide risk, no study has investigated the location of attachment within the ideation-to-action frameworks, with exception of study 29.

Our findings highlight that both anxious and avoidant attachment may lead to STB, but the mechanisms involved in this vulnerability development for each attachment orientation are still unclear. Mikulincer and Shaver (2016b), for example, propose that a suicide attempt could be the result of an anxiety-mediated hyperactivation of the attachment system, communicating the need for help and indicating that the person is struggling to cope with adversities and emotional pain. Levi-Belz et al. (study 24) is another example that demonstrates how avoidant and anxious attachment may increase one's vulnerability for a suicide attempt. These authors propose that the negative perception of others (internal working model of others), characteristic of high avoidant individuals, makes it precarious for them to disclose their emotions and thoughts to others. These attachment processes, in turn, reduce such individuals' ability to approach someone and ask for help in an emotional crisis. As a consequence, they are more likely to become detached,

lonely, and alienated, which increases the risk of suicidal behavior. Falgares et al. (study 8) offered a similar interpretation for their results finding that, having a negative view of others, individuals with higher levels of attachment avoidance may feel the need to be highly competent or nearly flawless at life tasks to maintain self-reliance rather than risk potential rejection. A suicide attempt could be a response to interpersonal rejection and/or to failing to reach these standards of perfectionism.

There was no evidence that high-quality studies had a different relationship between attachment and STB compared to low-quality studies. When evaluating the quality of the studies, we considered how attachment and STB are assessed. It is known, for instance, that adult attachment can be assessed both through self-administered scales (e.g., Arnsden and Greenberg, 1987; Griffin and Bartholomew, 1994) and clinical/research interviews (e.g., George et al., 1985; Hesse, 2008). These procedures are not only grounded in different approaches to attachment, they also can yield very different results which vary as a function of the nature of the assessment (e.g., an interview versus questionnaire, typological versus dimensional conceptualization). Although most studies assessed attachment and STB through self-administered measures, STB were most often assessed via clinical/research interviews when the participants were patients, and most studies, irrespective of sample characteristics, used self-administered scales to assess attachment. Therefore, it is difficult to say whether the assessment method differentially impacted the findings.

There are a number of clinical implications that arise from this review. Centrally, clinical assessment of patients should include an evaluation of their attachment relationships, their views of

themselves and of others, and, thus, their potential approach to coping with a mental health crisis involving suicidal ideation and attempts. Clinicians should employ both self-report measures and clinical interviews when possible (see Ravitz et al., 2010 for an extensive review of adult attachment measures) to examine the patient's beliefs and behaviors about close relationships. Such assessments may provide useful information about the patient's vulnerability associated with attachment insecurities as well as their coping strategies to deal with difficult life events and mental health crises. For example, if a patient tends to engage in anxious hyperactivation strategies (Mikulincer & Shaver, 2016d) and does not find the support needed from their attachment figures, feelings of thwarted belongingness may emerge and increase the likelihood of suicidal thoughts (as suggested by study 29). Such information may be useful when establishing a safety-planning intervention (Stanley & Brown, 2012), for instance. By being aware of a patient's attachment-related strategies (anxious hyperactivation or avoidant deactivating) during a crisis, both the clinician and the patient will be able to collaboratively plan a strategy that involves internal coping strategies and social support to mitigate against suicide risk. Such awareness will also provide clues about likely engagement with the therapeutic process. Some psychotherapeutic approaches that include attachment as a key construct, such as mentalization-based treatment (MBT; e.g., Bateman & Fonagy, 2013) and attachment-based family therapy (ABFT; e.g., Ewing, Diamond, & Levy, 2015) can provide clinicians a wide range of tools for assessment, formulation, and intervention.

The results of the present systematic review should be interpreted within the

context of its limitations. Conducting a meta-analysis of the studies included within this review was not feasible due to the level of heterogeneity across the studies regarding (1) the variety of conceptual frameworks of attachment and STB, (2) the diversity of instruments assessing attachment and suicidal thinking and behaviors, and (3) the heterogeneous populations participating in the studies. Therefore, we conducted a narrative review, which may be more susceptible to bias and subjectivity than a meta-analysis. We can also say little about whether the attachment-STB relationship differs across cultures because the vast majority (74%) of the participants in the 52 studies were from the United States. The attachment-STB relationship appears to hold irrespective of whether the samples were clinical or not. It is important to note, however, that we did not investigate publication and selection biases and we did not preregister a systematic review.

A significant challenge in integrating the attachment-STB research literature is the inconsistent use of terminology used to describe different patterns of attachment. This is, in part, because attachment theory comprises different conceptual approaches including Bowlby and Ainsworth's original conceptualization (e.g., Ainsworth, 1964; Ainsworth & Bowlby, 1991; Bowlby, 1988) and the contemporary frameworks on adult attachment (e.g., Bartholomew & Horowitz, 1991; Hazan & Shaver, 1987; Mikulincer & Shaver, 2016c). This inconsistency complicates the process of synthesizing the studies since the authors conceptualize attachment in such diverse ways, even though they are analyzing the same psychological phenomena. Equally, the term *insecure attachment* (e.g., study 35) limits our understanding of the specific attachment orientation that is being referred

to, since insecure attachment encapsulates a whole range of patterns including being anxious, dismissive, preoccupied, fearful, or disorganized. A similar concern can be extended to the various definitions of STB. Suicide risk, suicide tendencies, suicidal triggers, and suicidality have all been employed to describe suicidal ideation or behavior in a non-uniform manner. Studies assessing suicidal ideation and behaviors as a mixed concept, i.e., suicidality, may restrict the understanding of their relationship with attachment further, as suicidal ideation and suicide attempts are different phenomena and have unique psychological characteristics (O'Connor & Kirtley, 2018). Terms such as *suicidality* and *suicidal behaviors* are often defined in a vague manner (Silverman, 2006). A similar criticism can also be applied to uses of self-harm or self-injury to encompass both suicide attempt and non-suicidal self-injurious behaviors.

This review generates key research questions, such as (1) whether attachment plays a role in the differentiation between those who suffer from suicidal ideation and those who have a history of suicide attempt and (2) whether and how specific attachment dimensions might play different functions in this differentiation. It highlights (3) the need for empirical testing of what role attachment play in the theoretical understanding of suicidal behavior (Joiner, 2005; Klonsky & May, 2015; O'Connor, 2011); and lastly asks (4) What is the role of gender in the attachment-STB relationship?

CONCLUSION

The findings of this systematic review reveal that secure attachment seems to act as a protective factor against suicidal ideation and suicide attempt, and insecure dimensions of

attachment seem to function as vulnerability factors for STB. A synthesis of the studies' explanations of how attachment may be related to suicidal ideation and suicide attempts indicates that, in interaction with other personality factors, attachment moderates or mediates the relationship between current negative events and psychological states that, in turn, may lead to STB. Implications for treatment include targeting attachment-related coping strategies to mitigate vulnerability for STB.

AUTHOR NOTE

Tiago C. Zortea, Suicidal Behaviour Research Laboratory, Institute of Health and Wellbeing, University of Glasgow, Academic Centre, Gartnavel Royal Hospital, Glasgow, Scotland, United Kingdom

Cindy M. Gray, Institute of Health and Wellbeing, University of Glasgow, Glasgow, Scotland, United Kingdom

Rory C. O'Connor, Suicidal Behaviour Research Laboratory, Institute of Health and Wellbeing, University of Glasgow, Academic Centre, Gartnavel Royal Hospital, Glasgow, Scotland, United Kingdom

Correspondence concerning this article should be addressed to Tiago C. Zortea, Suicidal Behaviour Research Laboratory, Institute of Health and Wellbeing, University of Glasgow, Academic Centre, Gartnavel Royal Hospital, 1055 Great Western Road, Glasgow G12 0XH, Scotland, United Kingdom. Email: t.carlos-zortea.1@research.gla.ac.uk

FUNDING

This work was supported by a scholarship from the Brazilian National Council for

Scientific and Technological Development to the primary author under Grant 203245/2014-8.

SUPPLEMENTAL DATA

Supplemental data for this article can be accessed at [publisher's weblink](#).

ORCID

Tiago C. Zortea  <http://orcid.org/0000-0002-8451-0555>

Cindy M. Gray  <http://orcid.org/0000-0002-4295-6110>

Rory C. O'Connor  <http://orcid.org/0000-0002-3650-4994>

REFERENCES

- Aaltonen, K., Näätänen, P., Heikkinen, M., Koivisto, M., Baryshnikov, I., Karpov, B., ... Isometsä, E. (2016). Differences and similarities of risk factors for suicidal ideation and attempts among patients with depressive or bipolar disorders. *Journal of Affective Disorders, 193*, 318–330. doi:10.1016/j.jad.2015.12.033
- Adam, K. S. (1973). Childhood parental loss, suicidal ideation and suicidal behavior. In E. J. Anthony (Ed.), *The child in his family: The impact of disease and death* (pp. 275–297). New York, NY: Wiley.
- Adam, K. S. (1994). Suicidal behavior and attachment: A developmental model. In M. B. Sperling & W. H. Berman (Eds.), *Attachment in adults: Clinical and developmental perspectives* (1st ed., pp. 275–298). New York, NY: Guilford Press.
- Adam, K. S., Sheldon-Keller, A. E., & West, M. (1996). Attachment organization and history of suicidal behavior in clinical adolescents. *Journal of Consulting and Clinical Psychology, 64*(2), 264–272. doi:10.1037/0022-006X.64.2.264
- Addis, M., & Linehan, M. M. (1989). *Predicting suicidal behavior: Psychometric properties of the suicidal behaviors questionnaire*. In Poster presented at the annual meeting of the Association for the Advancement of Behavior Therapy, Washington, DC.
- Ainsworth, M. D. S. (1964). Patterns of attachment behavior shown by the infant in interaction with his mother. *Merrill-Palmer Quarterly of Behavior and Development, 10*(1), 51–58.
- Ainsworth, M. D. S., & Bowlby, J. (1991). An ethological approach to personality development. *American Psychologist, 46*(4), 333–341. doi:10.1037/0003-066X.46.4.333
- Ainsworth, M. D. S., Blehar, M. C., Waters, E., & Wall, S. (1978). *Patterns of attachment: A psychological study of the strange situation*. Hillsdale, NJ: Lawrence Erlbaum Associates.
- Armsden, G. C., & Greenberg, M. T. (1987). The inventory of parent and peer attachment: Individual differences and their relationship to psychological well-being in adolescence. *Journal of Youth and Adolescence, 16*(5), 427–454. doi:10.1007/BF02202939
- Baiocco, R., Ioverno, S., Lonigro, A., Baumgartner, E., & Laghi, F. (2015). Suicidal ideation among Italian and Spanish young adults: The role of sexual orientation. *Archives of Suicide Research, 19*(1), 75–88. doi:10.1080/13811118.2013.833150
- Bartholomew, K. (1990). Avoidance of intimacy: An attachment perspective. *Journal of Social and Personal Relationships, 7*(2), 147–178. doi:10.1177/0265407590072001
- Bartholomew, K., & Horowitz, L. M. (1991). Attachment styles among young adults: A test of a four-category model. *Journal of Personality and Social Psychology, 61*(2), 226–244. doi:10.1037/0022-3514.61.2.226
- Barzilay, S., & Apter, A. (2014). Psychological models of suicide. *Archives of Suicide Research, 18*(4), 295–241. doi:10.1080/13811118.2013.824825
- Bateman, A., & Fonagy, P. (2013). Mentalization-based treatment. *Psychoanalytic Inquiry, 33*(6), 595–613. doi:10.1080/07351690.2013.835170
- Bayam, G., Dilbaz, N., Bitlis, V., Holat, H., & Tüzer, T. (1995). İntihar davranış ile depresyon, ümitsizlik, intihar düşüncesi ilişkisi: İntihar Davranış Ölçeği geçerlik, güvenilirlik çalışması *Kriz Dergisi, 3*, 223–225.
- Beck, A. T., & Steer, R. A. (1991). *Manual for Beck Scale for suicide ideation*. New York, NY: Psychological Corporation.

Adult Attachment and Suicidal Thoughts and Behaviors

- Beck, A. T., & Steer, R. A. (1993). *Beck Scale for Suicide ideation manual*. San Antonio, TX: Psychological Corporation / Harcourt Brace & Company.
- Beck, A. T., Beck, R., & Kovacs, M. (1975). Classification of suicidal behaviors: I. Quantifying intent and medical lethality. *American Journal of Psychiatry*, *132*(3), 285–287. doi:10.1176/ajp.132.3.285
- Beck, A. T., Kovacs, M., & Weissman, A. (1979). Assessment of suicidal intention: The Scale for Suicide Ideation. *Journal of Consulting and Clinical Psychology*, *47*(2), 343–52. doi:10.1037/0022-006X.47.2.343
- Beck, A. T., Schuyler, D., & Herman, I. (1974). Development of suicidal intent scales. In A. T. Beck, H. L. Resnik, & D. J. Lettieri (Eds.), *The prediction of suicide* (pp. 45–56). Charles Press.
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the beck depression inventory-II*. San Antonio, TX: Psychological Corporation.
- Beck, A. T., Steer, R. A., & Ranieri, W. F. (1988). Scale for Suicide Ideation: Psychometric properties of a self-report version. *Journal of Clinical Psychology*, *44*(4), 499–505. doi:10.1002/1097-4679(198807)44:4 < 499::AID-JCLP2270440404 > 3.0.CO;2-6
- Beck, A. T., Ward, C. H., Mendelson, M., Mock, J., & Erbaugh, J. (1961). An inventory for measuring depression. *Archives of General Psychiatry*, *4*(6), 561–571. doi:10.1001/archpsyc.1961.01710120031004
- Bedi, R., Muller, R., & Classen, C. (2014). Cumulative risk for deliberate self-harm among treatment-seeking women with histories of childhood abuse. *Psychological Trauma: Theory Research Practice and Policy*, *6*(6), 600–609. doi:10.1037/a0033897
- Bowlby, J. (1988). *A secure base: Parent-child attachment and healthy human development*. New York, NY: Basic Books.
- Brennan, K. A., Clark, C. L., & Shaver, P. R. (1998). Self-report measurement of adult attachment: An integrative overview. In J. A. Simpson & W. S. Rholes (Eds.), *Attachment theory and close relationships* (pp. 46–76). New York, NY: Guilford Press.
- Brent, D. A., Perper, J. A., Moritz, G., Baugher, M., Roth, C., Balach, L., & Schweers, J. (1993). Stressful life events, psychopathology, and adolescent suicide: A case control Study. *Suicide & Life-Threatening Behavior*, *23*(3), 179–187. Retrieved from <http://search.ebscohost.com/login.aspx?direct=true&db=psyh&AN=1994-05893-001&site=ehost-live>
- Bretherton, I. (1992). The origins of attachment theory: John Bowlby and Mary Ainsworth. *Developmental Psychology*, *28*(5), 759–775. doi:10.1037/0012-1649.28.5.759
- Buelow, G., McClain, M., & McIntosh, I. (1996). A new measure for an important construct: The attachment and object relations inventory. *Journal of Personality Assessment*, *66*(3), 604–623. doi:10.1207/s15327752jpa6603_9
- Buelow, G., Schreiber, R., & Range, L. M. (2000). Attachment pattern, reasons for living, and suicide risk among college students. *Journal of College Counseling*, *3*(1), 29. doi:10.1002/j.2161-1882.2000.tb00161.x
- Canavarro, M. C., Dias, P., & Lima, V. (2006). A avaliação da vinculação do adulto: Uma revisão crítica a propósito da aplicação da Adult Attachment Scale-R (AAS-R) na população portuguesa. *Psicologia*, *20*, 155–186. doi:10.1007/s13398-014-0173-7.2
- Cero, I., & Sifers, S. K. (2013). Parenting behavior and the interpersonal-psychological theory of suicide: A mediated moderation analysis with adolescents. *Journal of Affective Disorders*, *150*(3), 987–992. doi:10.1016/j.jad.2013.05.025
- Chapman, A. L., Derbidge, C. M., Cooney, E., Hong, P. Y., & Linehan, M. M. (2009). Temperament as a prospective predictor of self-injury among patients with borderline personality disorder. *Journal of Personality Disorders*, *23*(2), 122–140. doi:10.1521/pedi.2009.23.2.122
- Cleare, S., Wetherall, K., Clark, A., Ryan, C., Kirtley, O., Smith, M., & O'Connor, R. (2018). Adverse childhood experiences and hospital-treated self-harm. *International Journal of Environmental Research and Public Health*, *15*(6), 1235. doi:10.3390/ijerph15061235
- Cloninger, C. R. (1992). *The temperament and character inventory*. Available from C. R. Cloninger. St. Louis, MO: Washington University School of Medicine, Department of Psychiatry.
- Cull, J. G., & Gill, W. S. (1982). *Suicide probability scale manual*. Los Angeles, CA: Western Psychological Services.
- Cruz, D., Narciso, I., Munoz, M., Pereira, C. R., & Sampaio, D. (2013). Adolescents and self-

- destructive behaviors: An exploratory analysis of family and individual correlates. *Behavioral Psychology/Psicologia Conductual*, 21(2), 271–288.
- Dilbaz, N., Holat, H., Bayam, G., & Tu?zer, T. (1995). Intihar düşüncesi ölçeğinin geçerlilik ve güvenilirliği. *Ulusal Psikiyatri Bilimler Kongresi Bilimsel Çalışma Kitabı*, 31, 40–41.
- D’Orío, B. M., Thompson, M. P., Lamis, D. A., Heron, S., Kaslow, N. J., D’Orío, B. M., ... Kaslow, N. J. (2015). Social support, attachment, and drug misuse in suicidal African American women. *Addiction Research & Theory*, 23(2), 170–176. doi:10.3109/16066359.2014.966096
- de Jong, M. L. (1992). Attachment, individuation, and risk of suicide in late adolescence. *Journal of Youth and Adolescence*, 21(3), 357–373. doi:10.1007/BF01537023
- DiFilippo, J. M., & Overholser, J. C. (2000). Suicidal ideation in adolescent psychiatric inpatients as associated with depression and attachment relationships. *Journal of Clinical Child Psychology*, 29(2), 155–166. doi:10.1207/S15374424jccp2902_2
- Ewing, E. S. K., Diamond, G., & Levy, S. (2015). Attachment-based family therapy for depressed and suicidal adolescents: Theory, clinical model and empirical support. *Attachment & Human Development*, 17(2), 136–156. doi:10.1080/14616734.2015.1006384
- Falgares, G., Marchetti, D., De Santis, S., Carrozzino, D., Kopala-Sibley, D. C., Fulcheri, M., & Verrocchio, M. C. (2017). Attachment styles and suicide-related behaviors in adolescence: The mediating role of self-criticism and dependency. *Frontiers in Psychiatry*, 8, 36. doi:10.3389/fpsyt.2017.00036
- Feeney, J. A., Noller, P., & Hanrahan, M. (1994). Assessing adult attachment. In M. B. Sperling & W. H. Berman (Eds.), *Attachment in adults: Clinical and developmental perspectives* (1st ed., pp. 128–152). New York, NY: Guilford Press.
- Fergusson, D. M., & Lynskey, M. T. (1995). Childhood circumstances, adolescent adjustment, and suicide attempts in a New Zealand birth cohort. *Journal of the American Academy of Child & Adolescent Psychiatry*, 34(5), 612–622. doi:10.1097/00004583-199505000-00013
- Fergusson, D. M., Woodward, L. J., & Horwood, L. J. (2000). Risk factors and life processes associated with the onset of suicidal behaviour during adolescence and early adulthood. *Psychological Medicine*, 30(1), 23–39. doi:10.1017/S003329179900135X
- Ferreira, J., & Castela, M. (1999). Questionário de Ideação Suicida (Q.I.S.). In M. R. Simões, M. M. Gonçalves, & L. S. Almeida (Eds.), *Testes e Provas Psicológicas em Portugal – Vol II* (pp. 123–130). Braga, Portugal: APPORT/SHO.
- Fraley, R. C., Heffernan, M. E., Vicary, A. M., & Brumbaugh, C. C. (2011). The experiences in close relationships-Relationship Structures Questionnaire: A method for assessing attachment orientations across relationships. *Psychological Assessment*, 23(3), 615–625. doi:10.1037/a0022898
- Fraley, R. C., Waller, N. G., & Brennan, K. A. (2000). An item-response theory analysis of self-report measures of adult attachment. *Journal of Personality and Social Psychology*, 78(2), 350–365. doi:10.1037/0022-3514.78.2.350
- Franklin, J. C., Ribeiro, J. D., Fox, K. R., Bentley, K. H., Kleiman, E. M., Huang, X., ... Nock, M. K. (2017). Risk factors for suicidal thoughts and behaviors: A meta-analysis of 50 years of research. *Psychological Bulletin*, 143(2), 187–232. doi:10.1037/bul0000084
- Garrison, C. Z., McKeown, R. E., Valois, R. F., & Vincent, M. L. (1993). Aggression, substance use, and suicidal behaviors in high school students. *American Journal of Public Health*, 83(2), 179–184. doi:10.2105/AJPH.83.2.179
- Genuis, M., Violato, C., Oddone-Paolucci, E., Robitaille, S., & McBride, D. (2000). Pilot study of the psychometric properties of the Adolescent Attachment Survey (AAS). In C. Violato, E. Oddone-Paolucci, & M. Genuis (Eds.), *The changing family and child development*. Aldershot, UK: Ashgate Publishing Limited.
- George, C., Kaplan, N., & Main, M. (1985). *The Berkeley adult attachment interview*. Berkeley, CA: University of California, Unpublished Manuscript.
- George, C., West, M., & Pettem, O. (1997). *The adult attachment projective. Unpublished attachment measure and coding manual*. Oakland, CA: Mills College.
- Gilreath, T. D., King, G., Graham, J. W., Flisher, A. J., & Lombard, C. (2009). Associations between maternal closeness, suicidal ideation,

- and risk behaviors in Cape Town. *European Child & Adolescent Psychiatry*, 18(3), 174–179. doi:10.1007/s00787-008-0718-4
- Glazebrook, K., Townsend, E., & Sayal, K. (2015). The role of attachment style in predicting repetition of adolescent self-harm: A longitudinal Study. *Suicide and Life-Threatening Behavior*, 45(6), 664–678. doi:10.1111/sltb.12159
- Goncalves, A. M., Sequeira, C. A. D., Duarte, J. C., & de Freitas, P. P. (2016). Suicidal ideation on higher education students: Influence of some psychosocial variables. *Archives of Psychiatric Nursing*, 30(2), 162–166. doi:10.1016/j.apnu.2015.08.005
- Gormley, B., & McNeil, D. E. (2010). Adult attachment orientations, depressive symptoms, anger, and self-directed aggression by psychiatric patients. *Cognitive Therapy and Research*, 34(3), 272–281. doi:10.1007/s10608-009-9267-5
- Griffin, D. W., & Bartholomew, K. (1994). The metaphysics of measurement: The case of adult attachment. In K. Bartholomew & D. Perlman (Eds.), *Advances in personal relationships: attachment processes in adulthood* (Vol. 5, pp. 17–52). London, UK: Jessica Kingsley Publishers.
- Grunebaum, M. F., Galfalvy, H. C., Mortenson, L. Y., Burke, A. K., Oquendo, M. A., & Mann, J. J. (2010). Attachment and social adjustment: Relationships to suicide attempt and major depressive episode in a prospective Study. *Journal of Affective Disorders*, 123(1-3), 123–130. doi:10.1016/j.jad.2009.09.010
- Hawton, K., Saunders, K. E., & O'Connor, R. C. (2012). Self-harm and suicide in adolescents. *The Lancet*, 379(9834), 2373–2382. doi:10.1016/S0140-6736(12)60322-5
- Hazan, C., & Shaver, P. (1987). Romantic love conceptualized as an attachment process. *Journal of Personality and Social Psychology*, 52(3), 511–524. doi:10.1037/0022-3514.52.3.511
- Hazan, C., & Shaver, P. R. (1994). Attachment as an organizational framework for research on close relationships. *Psychological Inquiry*, 5(1), 1–22. doi:10.1207/s15327965pli0501_1
- Hesse, E. (2008). The adult attachment interview: Protocol, method of analysis, and empirical studies. In J. Cassidy & P. R. Shaver (Eds.), *Handbook of attachment: Theory, research, and clinical applications* (2nd ed., pp. 552–598). New York, NY: Guilford Press.
- Heydari, A., Teymoori, A., & Nasiri, H. (2015). The effect of parent and peer attachment on suicidality: The mediation effect of self-control and anomie. *Community Mental Health Journal*, 51(3), 359–364. doi:10.1007/s10597-014-9809-5
- Ilardi, D. L., & Kaslow, N. J. (2009). Social difficulties influence group psychotherapy adherence in abused, suicidal African American women. *Journal of Clinical Psychology*, 65(12), 1300–1311. doi:10.1002/jclp.20628
- Joiner, T. (2005). *Why people die by suicide*. Cambridge, MA: Harvard University Press.
- Kerns, K. A., Schlegelmilch, A., Morgan, T. A., & Abraham, M. M. (2005). Assessing attachment in middle childhood. In K. A. Kerns & R. A. Richardson (Eds.), *Attachment in middle childhood* (1st ed., pp. 46–71). New York, NY: Guilford Press. Retrieved from http://books.google.com/books?hl=en&lr=&id=XX0_qY31iTUC&pgis=1
- Kessler, R. C., Borges, G., & Walters, E. E. (1999). Prevalence of and risk factors for lifetime suicide attempts in the National Comorbidity Survey. *Archives of General Psychiatry*, 56(7), 617–626. doi:10.1001/archpsyc.56.7.617
- Kidd, S., & Shahar, G. (2008). Resilience in homeless youth: The key role of self-esteem. *American Journal of Orthopsychiatry*, 78(2), 163–172. doi:10.1037/0002-9432.78.2.163
- Klonsky, E. D., & May, A. M. (2015). The Three-Step Theory (3ST): A new theory of suicide rooted in the “Ideation-to-Action” framework. *International Journal of Cognitive Therapy*, 8(2), 114–129. doi:10.1521/ijct.2015.8.2.114
- Klonsky, E. D., May, A. M., & Saffer, B. Y. (2016). Suicide, suicide attempts, and suicidal ideation. *Annual Review of Clinical Psychology*, 12(1), 307–330. doi:10.1146/annurev-clinpsy-021815-093204
- Kroenke, K., & Spitzer, R. L. (2002). The PHQ-9: A new depression diagnostic and severity measure. *Psychiatric Annals*, 32(9), 509–515. doi:10.3928/0048-5713-20020901-06
- Lessard, J. C., & Moretti, M. M. (1998). Suicidal ideation in an adolescent clinical sample: Attachment patterns and clinical implications. *Journal of Adolescence*, 21(4), 383–395. doi:10.1006/jado.1998.0169
- Levi-Belz, Y., Gvion, Y., Horesh, N., & Apter, A. (2013). Attachment patterns in medically serious suicide attempts: The mediating role of

- self-disclosure and loneliness. *Suicide and Life-Threatening Behavior*, 43(5), 511–522. doi:10.1111/sltb.12035
- Lewinsohn, P. M., Rohde, P., & Seeley, J. R. (1996). Adolescent suicidal ideation and attempts: Prevalence, risk factors, and clinical implications. *Clinical Psychology: Science and Practice*, 3(1), 25–46. doi:10.1111/j.1468-2850.1996.tb00056.x
- Linehan, M. M., & Nielsen, S. L. (1981). Assessment of suicide ideation and parasuicide: Hopelessness and social desirability. *Journal of Consulting and Clinical Psychology*, 49(5), 773–775. doi:10.1037/0022-006X.49.5.773
- Linehan, M. M., Comtois, K. A., Brown, M. Z., Heard, H. L., & Wagner, A. (2006). Suicide Attempt Self-Injury Interview (SASII): Development, reliability, and validity of a scale to assess suicide attempts and intentional self-injury. *Psychological Assessment*, 18(3), 303–312. doi:10.1037/1040-3590.18.3.303
- Liu, R. X. (2005). Parent-youth closeness and youth's suicidal ideation: The moderating effects of gender. *Stages of Adolescence, and Race or Ethnicity. Youth & Society*, 37(2), 145–175. doi:10.1177/0044118x04272290
- Livesley, W. J., Jang, K. L., & Vernon, P. A. (1998). Phenotypic and genetic structure of traits delineating personality disorder. *Archives of General Psychiatry*, 55(10), 941–948. doi:10.1001/archpsyc.55.10.941
- Lizardi, D., Grunebaum, M., Burke, A., Stanley, B., Mann, J., Harkavy-Friedman, J., & Oquendo, M. (2011). The effect of social adjustment and attachment style on suicidal behaviour. *Acta Psychiatrica Scandinavica*, 124(4), 295–300. doi:10.1111/j.1600-0447.2011.01724.x
- Maimon, D., & Kuhl, D. C. (2008). Social control and youth suicidality: Situating Durkheim's ideas in a multilevel framework. *American Sociological Review*, 73(6), 921–943. doi:10.1177/000312240807300603
- Main, M., Goldwyn, R., & Hesse, E. (2002). *Adult attachment scoring and classification system*. Unpublished manuscript. Berkeley, CA: University of California.
- Mann, J. J., McBride, P. A., Brown, R. P., Linnoila, M., Leon, A. C., DeMeo, M., ... Stanley, M. (1992). Relationship between central and peripheral serotonin indexes in depressed and suicidal psychiatric inpatients. *Archives of General Psychiatry*, 49(6), 442–446. doi:10.1001/archpsyc.1992.01820060022003
- Matos, P. M., & Costa, M. E. (2001). *Questionário de vinculação ao pai e à mãe. Manuscrito não publicado [The father and mother attachment questionnaire]*. Porto: Faculty of Psychology and Educational Sciences.
- Matos, P. M., Barbosa, S., & Costa, M. E. (2001). Avaliação da vinculação amorosa em adolescentes e jovens adultos: Construção de um instrumento e estudos de validação [The Romantic Attachment Questionnaire: Development and validation studies]. *Revista Oficial de La Asociación Iberoamericana de Diagnóstico Y Evaluación Psicológica*, 11, 93–109.
- Mikulincer, M., & Shaver, P. R. (2008). Adult attachment and affect regulation. In J. Cassidy & P. R. Shaver (Eds.), *Handbook of attachment: Theory, research and clinical applications* (2nd ed., pp. 503–531). New York, NY: The Guildford Press.
- Mikulincer, M., & Shaver, P. R. (2016a). An attachment perspective on interpersonal regulation. In M. Mikulincer & P. R. Shaver (Eds.), *Attachment in adulthood: Structure, dynamics, and change* (2nd ed., pp. 261–298). New York, NY: Guilford Press.
- Mikulincer, M., & Shaver, P. R. (2016b). Attachment bases of psychopathology. In M. Mikulincer & P. R. Shaver (Eds.), *Attachment in adulthood: Structure, dynamics, and change* (2nd ed., pp. 395–442). New York, NY: Guilford Press.
- Mikulincer, M., & Shaver, P. R. (2016c). *Attachment in adulthood: Structure, dynamics, and change* (2nd ed.). New York, NY: Guilford Press.
- Mikulincer, M., & Shaver, P. R. (2016d). A model of attachment-system functioning and dynamics in adulthood. In M. Mikulincer & P. R. Shaver (Eds.), *Attachment in adulthood: Structure, dynamics, and change* (2nd ed., pp. 395–442). New York, NY: Guilford Press.
- Miniati, M., Callari, A., & Pini, S. (2017). Adult attachment style and suicidality. *Psichiatria Danubina*, 29(3), 250–259. doi:10.24869/psyd.2017.250
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & The PRISMA Group. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement.

- PLoS Medicine*, 6(7), e1000097. doi:10.1371/journal.pmed1000097
- Muris, P., Mayer, B., & Meesters, C. (2000). Self-reported attachment style, anxiety, and depression in children. *Social Behavior and Personality*, 28(2), 157–162. doi:10.2224/sbp.2000.28.2.157
- Nagra, G. S., Lin, A., Upthegrove, R., G.S. N., & A, L. (2016). What bridges the gap between self-harm and suicidality? The role of forgiveness, resilience and attachment. *Psychiatry Research*, 241, 78–82. doi:10.1016/j.psychres.2016.04.103
- Nunes, F., & Mota, C. P. (2016). Parenting styles and suicidal ideation in adolescents: Mediating effect of attachment. *Journal of Child and Family Studies*, 26(3), 734–747. doi:10.1007/s10826-016-0611-6
- Nye, E. C., Katzman, J., Bell, J. B., Kilpatrick, J., Brainard, M., & Haaland, K. Y. (2009). Attachment representation and suicidal ideation in Vietnam combat veterans with posttraumatic stress disorder. *Archives of Suicide Research*, 13(2), 195–199. doi:10.1080/13811110902835213
- O'Connor, D. B., Ferguson, E., Green, J. A., O'carroll, R. E., & O'Connor, R. C. (2016). Cortisol levels and suicidal behavior: A meta-analysis. *Psychoneuroendocrinology*, 63, 370–379.
- O'Connor, R. C. (2011). Towards an integrated motivational-volitional model of suicidal behaviour. In R. C. O'Connor, S. Platt, & J. Gordon (Eds.), *International handbook of suicide prevention* (1st ed., pp. 181–198). Chichester, UK: John Wiley & Sons, Ltd.
- O'Connor, R. C., & Kirtley, O. J. (2018). The integrated motivational–volitional model of suicidal behaviour. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1754), 20170268. doi:10.1098/rstb.2017.0268
- O'Connor, R. C., & Nock, M. K. (2014). The psychology of suicidal behaviour. *The Lancet Psychiatry*, 1(1), 73–85. doi:10.1016/S2215-0366(14)70222-6
- Offord, D. R., Boyle, M. H., Szatmari, P., Rae Grant, N. I., Links, P. S., Cadman, D. T., ... Woodward, C. A. (1987). Ontario child health study II: Six-month prevalence of disorder and rates of service utilization. *Archives of General Psychiatry*, 44(9), 832–836. doi:10.1001/arch-psyc.1987.01800210084013
- Oquendo, M. A., Halberstam, B., & Mann, J. J. (2003). Risk factors for suicidal behavior: The utility and limitations of research instruments. In M. B. First (Ed.), *Standardized evaluation in clinical practice* (pp. 103–130). Washington, DC: American Psychiatric Publishing.
- Orbach, I., Milstein, I., Har-Even, D., Apter, A., Tiano, S., & Elizur, A. (1991). A multi-attitude suicide tendency scale for adolescents. *Psychological Assessment*, 3(3), 398–404. doi:10.1037/1040-3590.3.3.398
- Osman, A., Bagge, C. L., Gutierrez, P. M., Konick, L. C., Kopper, B. A., & Barrios, F. X. (2001). The Suicidal Behaviors Questionnaire-Revised (SBQ-R): Validation with clinical and nonclinical samples. *Assessment*, 8(4), 443–454. doi:10.1177/107319110100800409
- Ozbay, O., & Ozcan, Y. Z. (2006). A test of Hirschi's Social Bonding Theory: Juvenile delinquency in the high schools of Ankara, Turkey. *International Journal of Offender Therapy and Comparative Criminology*, 50(6), 711–726. doi:10.1177/0306624X05283525
- Özer, Ü., Yildirim, E. A., & Erkoç, Ş. N. (2015). Relationship of suicidal ideation and behavior to attachment style in patients with major depression. *Noro Psikiyatri Arsivi*, 52(3), 283–288. doi:10.5152/npa.2015.7459
- Palitsky, D., Mota, N., Affifi, T. O., Downs, A. C., & Sareen, J. (2013). The association between adult attachment style, mental disorders, and suicidality: Findings from a population-based study. *The Journal of Nervous and Mental Disease*, 201(7), 579–586. doi:10.1097/NMD.0b013e31829829ab
- Park, B. C. B., Kim, J. J., & Lester, D. (2011). Reasons for committing suicide in South Korean University students. *Suicidology Online*, 2, 11–16.
- Peter, T., Roberts, L. W., & Buzdugan, R. (2008). Suicidal ideation among Canadian Youth: A multivariate analysis. *Archives of Suicide Research*, 12(3), 263–275. doi:10.1080/13811110802100882
- Pfeffer, C. R. (1986). *The suicidal child* (1st ed.). New York, NY: The Guildford Press.
- Pisetsky, E. M., Wonderlich, S. A., Crosby, R. D., Peterson, C. B., Mitchell, J. E., Engel, S. G., ... Crow, S. J. (2015). Depression and personality traits associated with emotion dysregulation: Correlates of suicide attempts in women with bulimia nervosa. *European Eating Disorders Review*, 23(6), 537–544. doi:10.1002/erv.2401
- Posner, K., Brown, G. K., Stanley, B., Brent, D. A., Yershova, K. V., Oquendo, M. A., ... Mann, J. J. (2011). The Columbia-suicide

- severity rating scale: Initial validity and internal consistency findings from three multisite studies with adolescents and adults. *American Journal of Psychiatry*, 168(12), 1266–1277. doi:10.1176/appi.ajp.2011.10111704
- Raudino, A., Fergusson, D. M., & Horwood, L. J. (2013). The quality of parent/child relationships in adolescence is associated with poor adult psychosocial adjustment. *Journal of Adolescence*, 36(2), 331–340. doi:10.1016/j.adolescence.2012.12.002
- Ravitz, P., Maunder, R., Hunter, J., Sthankiya, B., & Lancee, W. (2010). Adult attachment measures: A 25-year review. *Journal of Psychosomatic Research*, 69(4), 419–432. doi:10.1016/j.jpsychores.2009.08.006
- Reich, W., Shayka, J. J., & Taibleson, C. (1991). *Manual to accompany the diagnostic interview for children and adolescents—revised*. St. Louis, MI: Washington University Press.
- Reynolds, W. M. (1991). Psychometric characteristics of the adult suicidal ideation Questionnaire in college students. *Journal of Personality Assessment*, 56(2), 289–307. doi:10.1207/s15327752jpa5602_9
- Richard, M. H., Fonagy, P., Smith, P. K., Wright, J. C., & Binney, V. (1998). *The adolescent separation anxiety interview*. Unpublished interview protocol. Goldsmiths College, University of London.
- Riggs, S. A., & Jacobvitz, D. (2002). Expectant parents' representations of early attachment relationships: Associations with mental health and family history. *Journal of Consulting and Clinical Psychology*, 70(1), 195–204. doi:10.1037//0022-006X.70.1.195
- Rodell, D. E., Benda, B. B., & Rodell, L. (2003). Suicidal thoughts among homeless alcohol and other drug abusers. *Alcoholism Treatment Quarterly*, 21(2), 57–74. doi:10.1300/J020v21n02_04
- Rodgers, R. F., van Leeuwen, N., Chabrol, H., & Leichsenring, F. (2011). An exploration of the role of defensive psychopathology in adolescent suicidal ideation and behavior. *Bulletin of the Menninger Clinic*, 75(3), 236–253. doi:10.1521/bumc.2011.75.3.236
- Ross, C. A., Heber, S., Norton, G. R., Anderson, D., Anderson, G., & Barchet, P. (1989). The dissociative disorders interview schedule: A structured interview. *Dissociation*, 2(3), 169–189.
- Rossotto, E., Yager, J., & Rorty, M. (1994). *Impulsive and self-harm behaviors among women with bulimia nervosa*. Paper presented at 6th Annual International Conference on Eating Disorders, New York.
- Rossouw, T. I., & Fonagy, P. (2012). Mentalization-based treatment for self-harm in adolescents: A randomized controlled trial. *Journal of the American Academy of Child & Adolescent Psychiatry*, 51(12), 1304–1313. doi:10.1016/j.jaac.2012.09.018
- Salzinger, S., Rosario, M., Feldman, R. S., & Ng-Mak, D. S. (2007). Adolescent suicidal behavior: Associations with preadolescent physical abuse and selected risk and protective factors. *Journal of the American Academy of Child & Adolescent Psychiatry*, 46(7), 859–866. doi:10.1097/chi.0b013e318054e702
- Scharfe, E., & Bartholomew, K. (1998). Do you remember? Recollections of adult attachment patterns. *Personal Relationships*, 5(2), 219–234. doi:10.1111/j.1475-6811.1998.tb00169.x
- Schotte, D. E., & Clum, G. A. (1982). Suicide ideation in a college population: A test of a model. *Journal of Consulting and Clinical Psychology*, 50(5), 690–696. doi:10.1037//0022-006X.50.5.690
- Sheftall, A. H., Mathias, C. W., Furr, R. M., & Dougherty, D. M. (2013). Adolescent attachment security, family functioning, and suicide attempts. *Attachment & Human Development*, 15(4), 368–383. doi:10.1080/14616734.2013.782649
- Sheftall, A., Schoppe-Sullivan, S., & Bridge, J. (2014). Insecure attachment and suicidal behavior in adolescents. *Crisis*, 35(6), 426–430. doi:10.1027/0227-5910/a000273
- Sibley, C. G., Fischer, R., & Liu, J. H. (2005). Reliability and validity of the revised experiences in close relationships (ECR-R) self-report measure of adult romantic attachment. *Personality and Social Psychology Bulletin*, 31(11), 1524–1536. doi:10.1177/0146167205276865
- Silverman, M. M. (2006). The language of suicidology. *Suicide and Life-Threatening Behavior*, 36(5), 519–532. doi:10.1521/suli.2006.36.5.519
- Silverman, M. M., Berman, A. L., Sanddal, N. D., O'Carroll, P. W., & Joiner, T. E. (2007). Rebuilding the Tower of Babel: A revised nomenclature for the study of suicide and suicidal behaviors part 2: Suicide-related ideations, communications, and behaviors. *Suicide and Life-Threatening Behavior*, 37(3), 264–277. doi:10.1521/suli.2007.37.3.264
- Simpson, J. A. (1990). Influence of attachment styles on romantic relationships. *Journal of*

Adult Attachment and Suicidal Thoughts and Behaviors

- Personality and Social Psychology*, 59(5), 971–980. doi:10.1037/0022-3514.59.5.971
- Simpson, J. A., & Rholes, W. S. (2012). Adult attachment orientations, stress, and romantic relationships. In P. Devine & A. Plant (Eds.), *Advances in experimental social psychology* (Vol. 45, pp. 179–328). Burlington, NJ: Academic Press. doi:10.1016/B978-0-12-394286-9.00006-8
- Smith, N. B., Mota, N., Tsai, J., Monteith, L., Harpaz-Rotem, I., Southwick, S. M., & Pietrzak, R. H. (2016). Nature and determinants of suicidal ideation among US Veterans: Results from the national health and resilience in veterans Study. *Journal of Affective Disorders*, 197, 66–73. doi:10.1016/j.jad.2016.02.069
- Stanley, B., & Brown, G. K. (2012). Safety planning intervention: A brief intervention to mitigate suicide risk. *Cognitive and Behavioral Practice*, 19(2), 256–264. doi:10.1016/j.cbpra.2011.01.001
- Steff, S., Morse, J., Yaggi, K., Reynolds, S., Reed, I., & Pilkonis, P. (2008). The role of attachment styles and interpersonal problems in suicide-related behaviors. *Suicide and Life-Threatening Behavior*, 38(5), 592–607. doi:10.1521/suli.2008.38.5.592
- Strang, S. P., & Orlofsky, J. L. (1990). Factors underlying suicidal ideation among college students: A test of Teicher and Jacobs' Model. *Journal of Adolescence*, 13(1), 39–52. doi:10.1016/0140-1971(90)90040-E
- Target, M., Fonagy, P., & Shmueli-Goetz, Y. (2003). Attachment representations in school-age children: The development of the Child Attachment Interview (CAI). *Journal of Child Psychotherapy*, 29, 171–186. doi:10.1080/0075417031000138433
- Turanovic, J. J., & Pratt, T. C. (2015). Longitudinal effects of violent victimization during adolescence on adverse outcomes in adulthood: A focus on prosocial attachments. *The Journal of Pediatrics*, 166(4), 1062–1069. doi:10.1016/j.jpeds.2014.12.059
- Turecki, G., & Brent, D. A. (2016). Suicide and suicidal behaviour. *The Lancet*, 387(10024), 1227–1239. doi:10.1016/S0140-6736(15)00234-2
- Van Leeuwen, N., Rodgers, R., Regner, I., & Chabrol, H. (2010). The role of acculturation in suicidal ideation among second-generation immigrant adolescents in France. *Transcultural Psychiatry*, 47(5), 812–832. doi:10.1177/1363461510382154
- Venta, A., Mellick, W., Schatte, D., & Sharp, C. (2014). Preliminary evidence that thoughts of thwarted belongingness mediate the relations between level of attachment insecurity and depression and suicide-related thoughts in inpatient adolescents. *Journal of Social and Clinical Psychology*, 33(5), 428–447. doi:10.1521/jscp.2014.33.5.428
- Violato, C., & Arato, J. (2004). Childhood attachment and adolescent suicide: A stepwise discriminant analysis in a case-comparison study. *Individual Differences Research*, 2(3), 162–168.
- Vrouva, I., Fonagy, P., Fearon, P. R. M., & Roussov, T. (2010). The risk-taking and self-harm inventory for adolescents: Development and psychometric evaluation. *Psychological Assessment*, 22(4), 852–865. doi:10.1037/a0020583
- West, M. L., Spreng, S. W., Rose, S. M., & Adam, K. S. (1999). Relationship between attachment-felt security and history of suicidal behaviours in clinical adolescents. *The Canadian Journal of Psychiatry*, 44(6), 578–582. doi:10.1177/070674379904400606
- World Health Organization. (2014). *Preventing suicide: A global imperative*. Geneva, Switzerland: WHO.
- Wrath, A. J., & Adams, G. C. (2018). Self-injurious behaviours and adult attachment: A review of the literature. *Archives of Suicide Research*, 1–24. doi:10.1080/13811118.2018.1486251
- Wright, J., Briggs, S., & Behringer, J. (2005). Attachment and the body in suicidal adolescents: A pilot Study. *Clinical Child Psychology and Psychiatry*, 10(4), 477–491. doi:10.1177/1359104505056310
- Yaseen, Z. S., Kopeykina, I., Gutkovich, Z., Bassirmia, A., Cohen, L. J., & Galynker, I. I. (2014). Predictive validity of the Suicide Trigger Scale (STS-3) for post-discharge suicide attempt in high-risk psychiatric inpatients. *PLoS One*, 9(1), e86768–9. doi:10.1371/journal.pone.0086768
- Zeyrek, E. Y. Y., Gençöz, F., Bergman, Y., Lester, D., Gencoz, F., Bergman, Y., & Lester, D. (2009). Suicidality, problem-solving skills, attachment style, and hopelessness in Turkish students. *Death Studies*, 33(9), 815–827. doi:10.1080/07481180903142407